

12/23/98
 3598 U.S. PTO

08/999,811, filed December 24, 1997, which is a CIP of Application Serial No. 08/207,550, filed on March 8, 1994, and is a CIP of Application Serial No. 08/465,968, filed on June 6, 1995

Please type a plus sign (+) inside this box →



UTILITY PATENT APPLICATION TRANSMITTAL <small>(Only for new nonprovisional applications under 37 CFR 1.53(b))</small>		Attny Docket No.	PF112P2D1	Total Pages	106
		First Named Inventor or Application Identifier			
		HU, et al.			
		Date	December 23, 1998		
APPLICATION ELEMENTS <small>See MPEP chapter 600 concerning utility patent application contents</small>		Assistant Commissioner for Patent Address to: Box Patent Application Washington, D.C. 20231			
1. <input checked="" type="checkbox"/> Fee Transmittal Form 2. <input checked="" type="checkbox"/> Specification [total pgs. 62] (preferred arrangement set forth below) - Descriptive title of the Invention - Cross References to Related Applications - Statement Regarding Fed Sponsored R&D - Reference to Microfiche Appendix - Background of the Invention - Brief Summary of the Invention - Brief Description of the Drawings (if filed) - Detailed Description - Claim(s) - Abstract of the Disclosure 3. <input checked="" type="checkbox"/> Drawing(s) Figs. 1A-13 Total Sheets: 19 4. <input checked="" type="checkbox"/> Oath or Declaration Total pages: 6 a. <input type="checkbox"/> Newly executed (original or copy) b. <input checked="" type="checkbox"/> Copy from a prior application (1.63(d)) (for con/div with Box 17 completed) i. <input type="checkbox"/> Deletion of Inventors (signed statement attached deleting inventor(s) named in prior app.) 5. <input checked="" type="checkbox"/> Incorporation by Reference (with Box 4b) The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.		6. <input type="checkbox"/> Microfiche Computer Program (appendix) 7. Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary) a. <input type="checkbox"/> Computer Readable Copy b. <input checked="" type="checkbox"/> Paper Copy (identical to computer copy) (14 pages) c. <input checked="" type="checkbox"/> Statement verifying identity of above copies		Accompanying Application Parts 8. <input type="checkbox"/> Assignment Papers (cover and document(s)) 9. <input type="checkbox"/> 37 CFR 3.73(b) Statement <input type="checkbox"/> Power of Attny 10. <input type="checkbox"/> English Translation Document 11. <input checked="" type="checkbox"/> Information Disclosure Statement/PTO-1449 <input type="checkbox"/> Copies of IDS Citations 12. <input type="checkbox"/> Preliminary Amendment 13. <input checked="" type="checkbox"/> Return Receipt Postcard (MPEP 503) (should be specifically itemized) 14. <input type="checkbox"/> Small Entity Statement(s) <input type="checkbox"/> Statement filed in prior application, status still proper and desired 15. <input type="checkbox"/> Certified copy of priority document(s) 16. <input type="checkbox"/> Other:	
17. If a CONTINUING APPLICATION, check appropriate box and supply the requisite information: <input type="checkbox"/> Continuation <input checked="" type="checkbox"/> Divisional <input type="checkbox"/> Continuation-in-Part (CIP) of prior application No.: 08/999,811, filed December 24, 1997, which is a CIP of Application Serial No. 08/207,550, filed on March 8, 1994, and is a CIP of Application Serial No. 08/465,968, filed on June 6, 1995					
18. CORRESPONDENCE ADDRESS <input type="checkbox"/> Customer Number or Bar Code Label or <input checked="" type="checkbox"/> Correspondence address below					
Name	Michele M. Wales, Esq., Attorney for Applicants Human Genome Sciences, Inc.				
Address	9410 Key West Avenue				
City	Rockville	State	MD	Zip Code	20850
Country	US	Telephone	301-309-8504	Fax	301-309-8439
Signature		Reg. No. P-43,975		Date	
Michele M. Wales				12/23/98	

3598 U.S. PTO

09/21/98

12/23/98

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Attorney Docket No.: PF112P2D1

Date: December 23, 1998

Assistant Commissioner for Patents
Box PATENT APPLICATION
Washington, D.C. 20231

Sir:

Transmitted herewith for filing under 37 C.F.R. §1.53 is the following patent application:

Applicants: HU et al.

Title of Invention: Vascular Endothelial Growth Factor 2

Including: -Specification (54 pgs);
Claims (7 pgs); and
Abstract (1 pg).

Drawings: Figs. 1A-13 (19 sheets)

Sequence Listing: 14 pages

PATENT APPLICATION FEE VALUE


The filing fee of the subject application being filed concurrently herewith has been calculated on the Preliminary Amendment also being filed concurrently herewith, as follows:

TYPE	NO. FILE	LESS	EXTRA	EXTRA RATE	FEE
Total Claims	32	-20	12	\$18.00 each	216.00
Independent	4	-3	1	\$78.00 each	78.00
Minimum Fee					760.00
Multiple Dependency Fee if applicable (\$260.00)					
Total Filing Fee					1,054.00

Priority of Application Serial No. 08/999,811, filed on December 24, 1997, Application Serial No. 08/207,550, filed on March 8, 1994, and Application Serial No. 08/465,968, filed on June 6, 1995 is hereby claimed under 35 U.S.C. §120.

Please charge the required fee to Deposit Account No. 08-3425. In addition, the Commissioner is hereby authorized to charge payment for any additional filing fees required under 37 C.F.R. 1.16 or credit any overpayment to Deposit Account No. 08-3425. A duplicate of this paper is attached.

Respectfully submitted,


Michele M. Wales (Reg. No. P-43,975)
Attorney for Applicants

Human Genome Sciences, Inc.
9410 Key West Avenue
Rockville, MD 20850
Telephone: (301) 610-5772

Enclosures

Vascular Endothelial Growth Factor 2

Cross-Reference to Related Applications

This application is a continuation-in-part of both U.S. Application No. 08/207,550, filed March 8, 1994 and U.S. Application No. 08/465,968, filed June 6, 1995. The content of all the aforesaid applications are relied upon and incorporated by reference in their entirety.

Background of the Invention

Field of the Invention

The present invention relates to newly identified polynucleotides, polypeptides encoded by such polynucleotides, the use of such polynucleotides and polypeptides, as well as the production of such polynucleotides and polypeptides. The polypeptides of the present invention have been identified as members of the vascular endothelial growth factor family. More particularly, the polypeptides of the present invention are human vascular endothelial growth factor 2 (VEGF2). The invention also relates to inhibiting the action of such polypeptides.

Related Art

The formation of new blood vessels, or angiogenesis, is essential for embryonic development, subsequent growth, and tissue repair. Angiogenesis is also an essential part of certain pathological conditions, such as neoplasia (*i.e.*, tumors and gliomas). Abnormal angiogenesis is associated with other diseases such as inflammation, rheumatoid arthritis, psoriasis, and diabetic retinopathy (Folkman, J. and Klagsbrun, M., *Science* 235:442-447(1987)).

Both acidic and basic fibroblast growth factor molecules are mitogens for endothelial cells and other cell types. Angiotropin and angiogenin can induce angiogenesis, although their functions are unclear (Folkman, J., *Cancer Medicine*,

Lea and Febiger Press, pp. 153-170 (1993)). A highly selective mitogen for vascular endothelial cells is vascular endothelial growth factor or VEGF (Ferrara, N. *et al.*, *Endocr. Rev.* 13:19-32 (1992)), which is also known as vascular permeability factor (VPF).

5 Vascular endothelial growth factor is a secreted angiogenic mitogen whose target cell specificity appears to be restricted to vascular endothelial cells. The murine VEGF gene has been characterized and its expression pattern in embryogenesis has been analyzed. A persistent expression of VEGF was observed in epithelial cells adjacent to fenestrated endothelium, *e.g.*, in choroid plexus and kidney glomeruli. The data was consistent with a role of VEGF as a multifunctional regulator of endothelial cell growth and differentiation (Breier, G. *et al.*, *Development* 114:521-532 (1992)).

10 VEGF shares sequence homology with human platelet-derived growth factors, PDGF α and PDGF β (Leung, D.W., *et al.*, *Science* 246:1306-1309, (1989)). The extent of homology is about 21% and 23%, respectively. Eight cysteine residues contributing to disulfide-bond formation are strictly conserved in these proteins. Although they are similar, there are specific differences between VEGF and PDGF. While PDGF is a major growth factor for connective tissue, VEGF is highly specific for endothelial cells. Alternatively spliced mRNAs have been identified for both VEGF, PLGF, and PDGF and these different splicing products differ in biological activity and in receptor-binding specificity. VEGF and PDGF function as homo-dimers or hetero-dimers and bind to receptors which elicit intrinsic tyrosine kinase activity following receptor dimerization.

20 VEGF has four different forms of 121, 165, 189 and 206 amino acids due to alternative splicing. VEGF121 and VEGF165 are soluble and are capable of promoting angiogenesis, whereas VEGF189 and VEGF206 are bound to heparin containing proteoglycans in the cell surface. The temporal and spatial expression of VEGF has been correlated with physiological proliferation of the blood vessels (Gajdusek, C.M., and Carbon, S.J., *Cell Physiol.* 139:570-579 (1989); McNeil,

25

30

P.L., *et al.*, *J. Cell. Biol.* 109:811-822 (1989)). Its high affinity binding sites are localized only on endothelial cells in tissue sections (Jakeman, L.B., *et al.*, *Clin. Invest.* 89:244-253 (1989)). The factor can be isolated from pituitary cells and several tumor cell lines, and has been implicated in some human gliomas (Plate, K.H., *Nature* 359:845-848 (1992)). Interestingly, expression of VEGF121 or VEGF165 confers on Chinese hamster ovary cells the ability to form tumors in nude mice (Ferrara, N. *et al.*, *J. Clin. Invest.* 91:160-170 (1993)). The inhibition of VEGF function by anti-VEGF monoclonal antibodies was shown to inhibit tumor growth in immune-deficient mice (Kim, K.J., *Nature* 362:841-844 (1993)). Further, a dominant-negative mutant of the VEGF receptor has been shown to inhibit growth of glioblastomas in mice.

Vascular permeability factor (VPF) has also been found to be responsible for persistent microvascular hyperpermeability to plasma proteins even after the cessation of injury, which is a characteristic feature of normal wound healing. This suggests that VPF is an important factor in wound healing. Brown, L.F. *et al.*, *J. Exp. Med.* 176:1375-1379 (1992).

The expression of VEGF is high in vascularized tissues, (*e.g.*, lung, heart, placenta and solid tumors) and correlates with angiogenesis both temporally and spatially. VEGF has also been shown to induce angiogenesis *in vivo*. Since angiogenesis is essential for the repair of normal tissues, especially vascular tissues, VEGF has been proposed for use in promoting vascular tissue repair (*e.g.*, in atherosclerosis).

U.S. Patent No. 5,073,492, issued December 17, 1991 to Chen *et al.*, discloses a method for synergistically enhancing endothelial cell growth in an appropriate environment which comprises adding to the environment, VEGF, effectors and serum-derived factor. Also, vascular endothelial cell growth factor C sub-unit DNA has been prepared by polymerase chain reaction techniques. The DNA encodes a protein that may exist as either a heterodimer or homodimer. The protein is a mammalian vascular endothelial cell mitogen and, as such, is useful

for the promotion of vascular development and repair, as disclosed in European Patent Application No. 92302750.2, published September 30, 1992.

Summary of the Invention

5 The polypeptides of the present invention have been putatively identified as a novel vascular endothelial growth factor based on amino acid sequence homology to human VEGF.

10 In accordance with one aspect of the present invention, there are provided novel mature polypeptides, as well as biologically active and diagnostically or therapeutically useful fragments, analogs, and derivatives thereof. The polypeptides of the present invention are of human origin.

15 In accordance with another aspect of the present invention, there are provided isolated nucleic acid molecules comprising polynucleotides encoding full length or truncated VEGF2 polypeptides having the amino acid sequences shown in SEQ ID NOS:2 or 4, respectively, or the amino acid sequences encoded by the cDNA clones deposited in bacterial hosts as ATCC Deposit Number 97149 on May 12, 1995 or ATCC Deposit Number 75698 on March 4, 1994.

The present invention also relates to biologically active and diagnostically or therapeutically useful fragments, analogs, and derivatives of VEGF2.

20 In accordance with still another aspect of the present invention, there are provided processes for producing such polypeptides by recombinant techniques comprising culturing recombinant prokaryotic and/or eukaryotic host cells, containing a nucleic acid sequence encoding a polypeptide of the present invention, under conditions promoting expression of said proteins and subsequent recovery of said proteins.

25 In accordance with yet a further aspect of the present invention, there are provided processes for utilizing such polypeptides, or polynucleotides encoding such polypeptides for therapeutic purposes, for example, to stimulate

angiogenesis, wound-healing, growth of damaged bone and tissue, and to promote vascular tissue repair.

In accordance with yet another aspect of the present invention, there are provided antibodies against such polypeptides and processes for producing such polypeptides.

In accordance with yet another aspect of the present invention, there are provided antagonists to such polypeptides, which may be used to inhibit the action of such polypeptides, for example, to prevent tumor angiogenesis and thus inhibit the growth of tumors, to treat diabetic retinopathy, inflammation, rheumatoid arthritis and psoriasis.

In accordance with another aspect of the present invention, there are provided nucleic acid probes comprising nucleic acid molecules of sufficient length to specifically hybridize to nucleic acid sequences of the present invention.

In accordance with another aspect of the present invention, there are provided methods of diagnosing diseases or a susceptibility to diseases related to mutations in nucleic acid sequences of the present invention and proteins encoded by such nucleic acid sequences.

In accordance with yet a further aspect of the present invention, there is provided a process for utilizing such polypeptides, or polynucleotides encoding such polypeptides, for *in vitro* purposes related to scientific research, synthesis of DNA and manufacture of DNA vectors.

These and other aspects of the present invention should be apparent to those skilled in the art from the teachings herein.

Brief Description of the Figures

The following drawings are illustrative of embodiments of the invention and are not meant to limit the scope of the invention as encompassed by the claims.

FIGS. 1A-1E show the full length nucleotide (SEQ ID NO:1) and the deduced amino acid (SEQ ID NO:2) sequence of VEGF2. The polypeptide comprises approximately 419 amino acid residues of which approximately 23 represent the leader sequence. The standard one letter abbreviations for amino acids are used. Sequencing was performed using the Model 373 Automated DNA Sequencer (Applied Biosystems, Inc.). Sequencing accuracy is predicted to be greater than 97%.

FIGS. 2A-2D show the nucleotide (SEQ ID NO:3) and the deduced amino acid (SEQ ID NO:4) sequence of a truncated, biologically active form of VEGF2. The polypeptide comprises approximately 350 amino acid residues of which approximately the first 24 amino acids represent the leader sequence.

FIGS. 3A-3B are an illustration of the amino acid sequence homology between PDGF α (SEQ ID NO:5), PDGF β (SEQ ID NO:6), VEGF (SEQ ID NO:7), and VEGF2 (SEQ ID NO:4). The boxed areas indicate the conserved sequences and the location of the eight conserved cysteine residues.

FIG. 4 shows, in table-form, the percent homology between PDGF α , PDGF β , VEGF, and VEGF2.

FIG. 5 shows the presence of VEGF2 mRNA in human breast tumor cell lines.

FIG. 6 depicts the results of a Northern blot analysis of VEGF2 in human adult tissues.

FIG. 7 shows a photograph of an SDS-PAGE gel after *in vitro* transcription, translation and electrophoresis of the polypeptide of the present invention. Lane 1: ^{14}C and rainbow M.W. marker; Lane 2: FGF control; Lane 3: VEGF2 produced by M13-reverse and forward primers; Lane 4: VEGF2 produced by M13 reverse and VEGF-F4 primers; Lane 5: VEGF2 produced by M13 reverse and VEGF-F5 primers.

FIGS. 8A and 8B depict photographs of SDS-PAGE gels. VEGF2 polypeptide was expressed in a baculovirus system consisting of Sf9 cells.

Protein from the medium and cytoplasm of cells were analyzed by SDS-PAGE under non-reducing (FIG.8A) and reducing (FIG.8B) conditions.

FIG. 9 depicts a photograph of an SDS-PAGE gel. The medium from Sf9 cells infected with a nucleic acid sequence of the present invention was precipitated. The resuspended precipitate was analyzed by SDS-PAGE and stained with coomassie brilliant blue.

FIG. 10 depicts a photograph of an SDS-PAGE gel. VEGF2 was purified from the medium supernatant and analyzed by SDS-PAGE in the presence or absence of the reducing agent β -mercaptoethanol and stained by coomassie brilliant blue.

FIG. 11 depicts reverse phase HPLC analysis of purified VEGF2 using a RP-300 column (0.21 x 3 cm, Applied Biosystems, Inc.). The column was equilibrated with 0.1% trifluoroacetic acid (Solvent A) and the proteins eluted with a 7.5 min gradient from 0 to 60% Solvent B, composed of acetonitrile containing 0.07% TFA. The protein elution was monitored by absorbance at 215 nm ("red" line) and 280 nm ("blue" line). The percentage of Solvent B is shown by the "green" line.

FIG. 12 is a bar graph illustrating the effect of partially-purified VEGF2 protein on the growth of vascular endothelial cells in comparison to basic fibroblast growth factor.

FIG. 13 is a bar graph illustrating the effect of purified VEGF2 protein on the growth of vascular endothelial cells.

Detailed Description of the Preferred Embodiments

In accordance with one aspect of the present invention, there are provided isolated nucleic acid molecules comprising a polynucleotide encoding a VEGF2 polypeptide having the deduced amino acid sequence of Figure 1 (SEQ ID NO:2), which was determined by sequencing a cloned cDNA. The nucleotide sequence shown in SEQ ID NO:1 was obtained by sequencing a cDNA clone, which was

deposited on May 12, 1995 at the American Type Tissue Collection (ATCC),
12301 Park Lawn Drive, Rockville, Maryland 20852, and given ATCC Deposit
No. 97149.

5 In accordance with another aspect of the present invention, there are
provided isolated nucleic acid molecules comprising a polynucleotide encoding
a truncated VEGF2 polypeptide having the deduced amino acid sequence of
Figure 2 (SEQ ID NO:4), which was determined by sequencing a cloned cDNA.
The nucleotide sequence shown in SEQ ID NO:3 was obtained by sequencing a
cDNA clone, which was deposited on March 4, 1994 at the American Type
10 Tissue Collection (ATCC), 12301 Park Lawn Drive, Rockville, Maryland 20852,
and given ATCC Deposit Number 75698.

15 Unless otherwise indicated, all nucleotide sequences determined by
sequencing a DNA molecule herein were determined using an automated DNA
sequencer (such as the Model 373 from Applied Biosystems, Inc.), and all amino
acid sequences of polypeptides encoded by DNA molecules determined herein
were predicted by translation of a DNA sequence determined as above.
Therefore, as is known in the art for any DNA sequence determined by this
automated approach, any nucleotide sequence determined herein may contain
some errors. Nucleotide sequences determined by automation are typically at
20 least about 90% identical, more typically at least about 95% to at least about
99.9% identical to the actual nucleotide sequence of the sequenced DNA
molecule. The actual sequence can be more precisely determined by other
approaches including manual DNA sequencing methods well known in the art.
As is also known in the art, a single insertion or deletion in a determined
25 nucleotide sequence compared to the actual sequence will cause a frame shift in
translation of the nucleotide sequence such that the predicted amino acid
sequence encoded by a determined nucleotide sequence will be completely
different from the amino acid sequence actually encoded by the sequenced DNA
molecule, beginning at the point of such an insertion or deletion.

A polynucleotide encoding a polypeptide of the present invention may be obtained from early stage human embryo (week 8 to 9) osteoclastomas, adult heart or several breast cancer cell lines. The polynucleotide of this invention was discovered in a cDNA library derived from early stage human embryo week 9. It is structurally related to the VEGF/PDGF family. It contains an open reading frame encoding a protein of about 419 amino acid residues of which approximately the first 23 amino acid residues are the putative leader sequence such that the mature protein comprises 396 amino acids, and which protein exhibits the highest amino acid sequence homology to human vascular endothelial growth factor (30% identity), followed by PDGF α (24%) and PDGF β (22%). (See FIG. 4). It is particularly important that all eight cysteines are conserved within all four members of the family (see boxed areas of FIG. 3). In addition, the signature for the PDGF/VEGF family, PXC VXXXRCXGCCN, (SEQ ID NO: 8) is conserved in VEGF2 (see FIG. 3). The homology between VEGF2, VEGF and the two PDGFs is at the protein sequence level. No nucleotide sequence homology can be detected, and therefore, it would be difficult to isolate the VEGF2 through simple approaches such as low stringency hybridization.

The VEGF2 polypeptide of the present invention is meant to include the full length polypeptide and polynucleotide sequence which encodes for any leader sequences and for active fragments of the full length polypeptide. Active fragments are meant to include any portions of the full length amino acid sequence which have less than the full 419 amino acids of the full length amino acid sequence as shown in SEQ ID NO:2, but still contain the eight cysteine residues shown conserved in Figure 3 and that still have VEGF2 activity.

There are at least two alternatively spliced VEGF2 mRNA sequences present in normal tissues. The two bands in FIG. 7, lane 5 indicate the presence of the alternatively spliced mRNA encoding the VEGF2 polypeptide of the present invention.

5 The polynucleotide of the present invention may be in the form of RNA or in the form of DNA, which DNA includes cDNA, genomic DNA, and synthetic DNA. The DNA may be double-stranded or single-stranded, and if single stranded may be the coding strand or non-coding (anti-sense) strand. The coding sequence which encodes the mature polypeptide may be identical to the coding sequence shown in FIG. 1 or FIG. 2, or that of the deposited clones, or may be a different coding sequence which, as a result of the redundancy or degeneracy of the genetic code, encodes the same, mature polypeptide as the DNA of FIG. 1, FIG. 2, or the deposited cDNAs.

10 The polynucleotide which encodes for the mature polypeptide of FIG. 1 or FIG. 2 or for the mature polypeptides encoded by the deposited cDNAs may include: only the coding sequence for the mature polypeptide; the coding sequence for the mature polypeptide and additional coding sequences such as a leader or secretory sequence or a proprotein sequence; the coding sequence for the mature polypeptide (and optionally additional coding sequences) and non-coding sequences, such as introns or non-coding sequence 5' and/or 3' of the coding sequence for the mature polypeptide.

15 Thus, the term "polynucleotide encoding a polypeptide" encompasses a polynucleotide which includes only coding sequences for the polypeptide as well as a polynucleotide which includes additional coding and/or non-coding sequences.

20 The present invention further relates to variants of the hereinabove described polynucleotides which encode for fragments, analogs, and derivatives of the polypeptide having the deduced amino acid sequence of FIGS. 1 or 2, or the polypeptide encoded by the cDNA of the deposited clones. The variant of the polynucleotide may be a naturally occurring allelic variant of the polynucleotide or a non-naturally occurring variant of the polynucleotide.

25 Thus, the present invention includes polynucleotides encoding the same mature polypeptide as shown in FIGS. 1 or 2 or the same mature polypeptide encoded by the cDNA of the deposited clones as well as variants of such

30

polynucleotides which variants encode for a fragment, derivative, or analog of the polypeptides of FIGS. 1 or 2, or the polypeptide encoded by the cDNA of the deposited clones. Such nucleotide variants include deletion variants, substitution variants, and addition or insertion variants.

5 As hereinabove indicated, the polynucleotide may have a coding sequence which is a naturally occurring allelic variant of the coding sequence shown in FIGS. 1 or 2, or of the coding sequence of the deposited clones. As known in the art, an allelic variant is an alternate form of a polynucleotide sequence which have a substitution, deletion or addition of one or more nucleotides, which does not substantially alter the function of the encoded polypeptide.

10 The present invention also includes polynucleotides, wherein the coding sequence for the mature polypeptide may be fused in the same reading frame to a polynucleotide which aids in expression and secretion of a polypeptide from a host cell, for example, a leader sequence which functions as a secretory sequence for controlling transport of a polypeptide from the cell. The polypeptide having a leader sequence is a preprotein and may have the leader sequence cleaved by the host cell to form the mature form of the polypeptide. The polynucleotides may also encode for a proprotein which is the mature protein plus additional 5' amino acid residues. A mature protein having a prosequence is a proprotein and is an inactive form of the protein. Once the prosequence is cleaved an active mature protein remains.

15 Thus, for example, the polynucleotide of the present invention may encode for a mature protein, or for a protein having a prosequence or for a protein having both a prosequence and presequence (leader sequence).

20 The polynucleotides of the present invention may also have the coding sequence fused in frame to a marker sequence which allows for purification of the polypeptide of the present invention. The marker sequence may be a hexahistidine tag supplied by a pQE-9 vector to provide for purification of the mature polypeptide fused to the marker in the case of a bacterial host, or, for example, the marker sequence may be a hemagglutinin (HA) tag when a mammalian host,

e.g. COS-7 cells, is used. The HA tag corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson, I., *et al.*, *Cell* 37:767 (1984)).

Further embodiments of the invention include isolated nucleic acid molecules comprising a polynucleotide having a nucleotide sequence at least 95% identical, and more preferably at least 96%, 97%, 98% or 99% identical to (a) a nucleotide sequence encoding the polypeptide having the amino acid sequence in SEQ ID NO:2; (b) a nucleotide sequence encoding the polypeptide having the amino acid sequence in SEQ ID NO:2, but lacking the N-terminal methionine; (c) a nucleotide sequence encoding the polypeptide having the amino acid sequence at positions from about 1 to about 396 in SEQ ID NO:2; (d) a nucleotide sequence encoding the polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No.97149; (e) a nucleotide sequence encoding the mature VEGF2 polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No.97149; or (f) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), (c), (d), or (e).

Further embodiments of the invention include isolated nucleic acid molecules comprising a polynucleotide having a nucleotide sequence at least 95% identical, and more preferably at least 96%, 97%, 98% or 99% identical to (a) a nucleotide sequence encoding the polypeptide having the amino acid sequence in SEQ ID NO:4; (b) a nucleotide sequence encoding the polypeptide having the amino acid sequence in SEQ ID NO:4, but lacking the N-terminal methionine; (c) a nucleotide sequence encoding the polypeptide having the amino acid sequence at positions from about 1 to about 326 in SEQ ID NO:4; (d) a nucleotide sequence encoding the polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No.75698; (e) a nucleotide sequence encoding the mature VEGF2 polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No.75698; or (f) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), (c), (d), or (e).

By a polynucleotide having a nucleotide sequence at least, for example, 95% "identical" to a reference nucleotide sequence encoding a VEGF2 polypeptide is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence encoding the VEGF2 polypeptide. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular nucleic acid molecule is at least 95%, 96%, 97%, 98% or 99% identical to, for instance, the nucleotide sequence shown in SEQ ID NOS:1 or 3, or to the nucleotides sequence of the deposited cDNA clone(s) can be determined conventionally using known computer programs such as the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711). Bestfit uses the local homology algorithm of Smith and Waterman, *Advances in Applied Mathematics* 2: 482-489 (1981), to find the best segment of homology between two sequences. When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference nucleotide sequence and that gaps in homology of up to 5% of the total number of nucleotides in the reference sequence are allowed.

The term "gene" means the segment of DNA involved in producing a polypeptide chain; it includes regions preceding and following the coding region (leader and trailer) as well as intervening sequences (introns) between individual coding segments (exons).

5 The present invention is further directed to fragments of the isolated nucleic acid molecules described herein. By a fragment of an isolated nucleic acid molecule having the nucleotide sequence of the deposited cDNA(s) or the nucleotide sequence shown in SEQ ID NO:1 or SEQ ID NO:3 is intended fragments at least about 15 nt, and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably, at least about 40 nt in length which are useful as diagnostic probes and primers as discussed herein. Of course, larger fragments 50-1500 nt in length are also useful according to the present invention as are fragments corresponding to most, if not all, of the nucleotide sequence of the deposited cDNA(s) or as shown in SEQ ID NO:1 or SEQ ID NO:3. By a fragment at least 20 nt in length, for example, is intended fragments which include 20 or more contiguous bases from the nucleotide sequence of the deposited cDNA(s) or the nucleotide sequence as shown in SEQ ID NOS:1 or 3.

10
15
20 Fragments of the full length gene of the present invention may be used as a hybridization probe for a cDNA library to isolate the full length cDNA and to isolate other cDNAs which have a high sequence similarity to the gene or similar biological activity. Probes of this type preferably have at least 30 bases and may contain, for example, 50 or more bases. The probe may also be used to identify a cDNA clone corresponding to a full length transcript and a genomic clone or clones that contain the complete gene including regulatory and promotor regions, exons, and introns. An example of a screen comprises isolating the coding region of the gene by using the known DNA sequence to synthesize an oligonucleotide probe. Labeled oligonucleotides having a sequence complementary to that of the gene of the present invention are used to screen a library of human cDNA,

25

genomic DNA or mRNA to determine which members of the library the probe hybridizes to.

The present invention further relates to isolated nucleic acid molecules comprising polynucleotides which hybridize under stringent hybridization conditions to a portion of the polynucleotides in nucleic acid molecules of the invention described above, for instance, the cDNA clone(s) contained in ATCC Deposit Nos. 97149 or 75698. By "stringent hybridization conditions" is intended overnight incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (150 mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 g/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C.

By a polynucleotide which hybridizes to a "portion" of a polynucleotide is intended a polynucleotide (either DNA or RNA) hybridizing to at least about 15 nucleotides (nt), and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably about 30-70 nt of the reference polynucleotide. These are useful as diagnostic probes and primers as discussed above and in more detail below.

By a portion of a polynucleotide of "at least 20 nt in length," for example, is intended 20 or more contiguous nucleotides from the nucleotide sequence of the reference polynucleotide (e.g., the deposited cDNA or the nucleotide sequence as shown in SEQ ID NO:1). Of course, a polynucleotide which hybridizes only to a poly A sequence (such as the 3' terminal poly(A) tract of the VEGF2 cDNA shown in SEQ ID NOS:1 or 3), or to a complementary stretch of T (or U) residues, would not be included in a polynucleotide of the invention used to hybridize to a portion of a nucleic acid of the invention, since such a polynucleotide would hybridize to any nucleic acid molecule containing a poly (A) stretch or the complement thereof (e.g., practically any double-stranded cDNA clone).

The present application is directed to nucleic acid molecules at least 95%, 96%, 97%, 98% or 99% identical to the nucleic acid sequence shown in SEQ ID NOS:1 or 3 or to the nucleic acid sequence of the deposited cDNA(s), irrespective

of whether they encode a polypeptide having VEGF2 activity. This is because even where a particular nucleic acid molecule does not encode a polypeptide having VEGF2 activity, one of skill in the art would still know how to use the nucleic acid molecule, for instance, as a hybridization probe or a polymerase chain reaction (PCR) primer. Uses of the nucleic acid molecules of the present invention that do not encode a polypeptide having VEGF2 activity include, *inter alia*, (1) isolating the VEGF2 gene or allelic variants thereof in a cDNA library; (2) *in situ* hybridization (*e.g.*, "FISH") to metaphase chromosomal spreads to provide precise chromosomal location of the VEGF2 gene, as described in Verma *et al.*, *Human Chromosomes: A Manual of Basic Techniques*, Pergamon Press, New York (1988); and Northern Blot analysis for detecting VEGF2 mRNA expression in specific tissues.

Preferred, however, are nucleic acid molecules having sequences at least 95%, 96%, 97%, 98% or 99% identical to a nucleic acid sequence shown in SEQ ID NOS:1 or 3 or to a nucleic acid sequence of the deposited cDNA(s) which do, in fact, encode a polypeptide having VEGF2 protein activity. By "a polypeptide having VEGF2 activity" is intended polypeptides exhibiting VEGF2 activity in a particular biological assay. For example, VEGF2 protein activity can be measured using, for example, mitogenic assays and endothelial cell migration assays. See, *e.g.*, Olofsson *et al.*, *Proc. Natl. Acad. Sci. USA* 93:2576-2581 (1996) and Joukov *et al.*, *EMBO J.* 5:290-298 (1996).

Of course, due to the degeneracy of the genetic code, one of ordinary skill in the art will immediately recognize that a large number of the nucleic acid molecules having a sequence at least 90%, 95%, 96%, 97%, 98%, or 99% identical to a nucleic acid sequence of the deposited cDNA(s) or the nucleic acid sequence shown in SEQ ID NO:1 or SEQ ID NO:3 will encode a polypeptide "having VEGF2 protein activity." In fact, since degenerate variants of these nucleotide sequences all encode the same polypeptide, this will be clear to the skilled artisan even without performing the above described comparison assay. It will be further recognized in the art that, for such nucleic acid molecules that

are not degenerate variants, a reasonable number will also encode a polypeptide having VEGF2 protein activity. This is because the skilled artisan is fully aware of amino acid substitutions that are either less likely or not likely to significantly effect protein function (*e.g.*, replacing one aliphatic amino acid with a second aliphatic amino acid).

For example, guidance concerning how to make phenotypically silent amino acid substitutions is provided in Bowie, J. U. *et al.*, "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," *Science* 247:1306-1310 (1990), wherein the authors indicate that proteins are surprisingly tolerant of amino acid substitutions.

Thus, the present invention is directed to polynucleotides having at least a 70% identity, preferably at least 90% and more preferably at least a 95% identity to a polynucleotide which encodes the polypeptides of SEQ ID NOS:2 or 4, as well as fragments thereof, which fragments have at least 30 bases and preferably at least 50 bases and to polypeptides encoded by such polynucleotides.

The deposit(s) referred to herein will be maintained under the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the purposes of Patent Procedure. These deposits are provided merely as a convenience and are not an admission that a deposit is required under 35 U.S.C. § 112. The sequence of the polynucleotides contained in the deposited materials, as well as the amino acid sequence of the polypeptides encoded thereby, are incorporated herein by reference and are controlling in the event of any conflict with the description of sequences herein. A license may be required to make, use or sell the deposited materials, and no such license is hereby granted.

The present invention further relates to polypeptides which have the deduced amino acid sequence of FIGS. 1 or 2, or which has the amino acid sequence encoded by the deposited cDNAs, as well as fragments, analogs, and derivatives of such polypeptides.

The terms "fragment," "derivative" and "analog" when referring to the polypeptide of FIGS. 1 or 2 or that encoded by the deposited cDNA, means a

polypeptide which retains the conserved motif of VEGF proteins as shown in FIG. 3 and essentially the same biological function or activity.

The polypeptides of the present invention may be recombinant polypeptides, natural polypeptides, or synthetic polypeptides, preferably recombinant polypeptides.

It will be recognized in the art that some amino acid sequences of the VEGF2 polypeptide can be varied without significant effect of the structure or function of the protein. If such differences in sequence are contemplated, it should be remembered that there will be critical areas on the protein which determine activity.

Thus, the invention further includes variations of the VEGF2 polypeptide which show substantial VEGF2 polypeptide activity or which include regions of VEGF2 protein such as the protein portions discussed below. Such mutants include deletions, insertions, inversions, repeats, and type substitutions. As indicated above, guidance concerning which amino acid changes are likely to be phenotypically silent can be found in Bowie, J.U., *et al.*, "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," *Science* 247:1306-1310 (1990).

Thus, the fragments, derivatives, or analogs of the polypeptides of FIGS. 1 or 2, or that encoded by the deposited cDNAs may be: (i) one in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code; or (ii) one in which one or more of the amino acid residues includes a substituent group; or (iii) one in which the mature polypeptide is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol); or (iv) one in which the additional amino acids are fused to the mature polypeptide, such as a leader or secretory sequence or a sequence which is employed for purification of the mature polypeptide or a proprotein sequence; or (v) one in which comprises fewer amino

acid residues shown in SEQ ID NOS: 2 or 4, and retains the conserved motif and yet still retains activity characteristics of the VEGF family of polypeptides. Such fragments, derivatives, and analogs are deemed to be within the scope of those skilled in the art from the teachings herein.

5 Of particular interest are substitutions of charged amino acids with another charged amino acid and with neutral or negatively charged amino acids. The latter results in proteins with reduced positive charge to improve the characteristics of the VEGF2 protein. The prevention of aggregation is highly desirable. Aggregation of proteins not only results in a loss of activity but can also be problematic when preparing pharmaceutical formulations, because they can be immunogenic. (Pinckard *et al.*, *Clin. Exp. Immunol.* 2:331-340 (1967); Robbins *et al.*, *Diabetes* 36:838-845 (1987); Cleland *et al.* *Crit. Rev. Therapeutic Drug Carrier Systems* 10:307-377 (1993)).

10 The replacement of amino acids can also change the selectivity of binding to cell surface receptors. Ostade *et al.*, *Nature* 361:266-268 (1993) describes certain mutations resulting in selective binding of TNF- α to only one of the two known types of TNF receptors. Thus, the VEGF2 of the present invention may include one or more amino acid substitutions, deletions or additions, either from natural mutations or human manipulation.

15 As indicated, changes are preferably of a minor nature, such as conservative amino acid substitutions that do not significantly affect the folding or activity of the protein (see Table 1).

20

TABLE 1. Conservative Amino Acid Substitutions

Aromatic	Phenylalanine Tryptophan Tyrosine
Hydrophobic	Leucine Isoleucine Valine
Polar	Glutamine Asparagine
Basic	Arginine Lysine Histidine
Acidic	Aspartic Acid Glutamic Acid
Small	Alanine Serine Threonine Methionine Glycine

Amino acids in the VEGF2 protein of the present invention that are essential for function can be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, *Science* 244:1081-1085 (1989)). The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for biological activity such as receptor binding or *in vitro*, or *in vitro* proliferative activity. Sites that are critical for ligand-receptor binding can also be determined by structural analysis such as crystallization, nuclear magnetic resonance or photoaffinity labeling (Smith *et al.*, *J. Mol. Biol.* 224:899-904 (1992) and de Vos *et al.* *Science* 255:306-312 (1992)).

The polypeptides and polynucleotides of the present invention are preferably provided in an isolated form, and preferably are purified to homogeneity.

The term "isolated" means that the material is removed from its original environment (*e.g.*, the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide or polypeptide present in a living

animal is not isolated, but the same polynucleotide or DNA or polypeptide, separated from some or all of the coexisting materials in the natural system, is isolated. Such polynucleotide could be part of a vector and/or such polynucleotide or polypeptide could be part of a composition, and still be isolated in that such vector or composition is not part of its natural environment.

The polypeptides of the present invention include the polypeptides of SEQ ID NOS:2 and 4 (in particular the mature polypeptide) as well as polypeptides which have at least 70% similarity (preferably at least 70% identity) to the polypeptides of SEQ ID NOS:2 and 4, and more preferably at least 90% similarity (more preferably at least 95% identity) to the polypeptides of SEQ ID NOS:2 and 4, and still more preferably at least 95% similarity (still more preferably at least 90% identity) to the polypeptides of SEQ ID NOS:2 and 4 and also include portions of such polypeptides with such portion of the polypeptide generally containing at least 30 amino acids and more preferably at least 50 amino acids.

As known in the art "similarity" between two polypeptides is determined by comparing the amino acid sequence and its conserved amino acid substitutes of one polypeptide to the sequence of a second polypeptide.

Fragments or portions of the polypeptides of the present invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, the fragments may be employed as intermediates for producing the full-length polypeptides. Fragments or portions of the polynucleotides of the present invention may be used to synthesize full-length polynucleotides of the present invention.

The present invention also relates to recombinant vectors, which include the isolated nucleic acid molecules of the present invention, and to host cells containing the recombinant vectors, as well as to methods of making such vectors and host cells and for using them for production of VEGF2 polypeptides or peptides by recombinant techniques.

Host cells are genetically engineered (transduced, transformed, or transfected) with the vectors of this invention which may be, for example, a

cloning vector or an expression vector. The vector may be, for example, in the form of a plasmid, a viral particle, a phage, etc. The engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants, or amplifying the VEGF2 genes of the invention. The culture conditions, such as temperature, pH and the like, are those previously used with the host cell selected for expression, and will be apparent to the skilled artisan.

The polynucleotides of the present invention may be employed for producing polypeptides by recombinant techniques. Thus, for example, the polynucleotide sequence may be included in any one of a variety of expression vectors for expressing a polypeptide. Such vectors include chromosomal, nonchromosomal and synthetic DNA sequences, *e.g.*, derivatives of SV40; bacterial plasmids; phage DNA; yeast plasmids; vectors derived from combinations of plasmids and phage DNA, viral DNA such as vaccinia, adenovirus, fowl pox virus, and pseudorabies. However, any other plasmid or vector may be used so long as it is replicable and viable in the host.

The appropriate DNA sequence may be inserted into the vector by a variety of procedures. In general, the DNA sequence is inserted into an appropriate restriction endonuclease site(s) by procedures known in the art. Such procedures and others are deemed to be within the scope of those skilled in the art.

The DNA sequence in the expression vector is operatively linked to an appropriate expression control sequence(s) (promoter) to direct mRNA synthesis. As representative examples of such promoters, there may be mentioned: LTR or SV40 promoter, the *E. coli. lac* or *trp*, the phage lambda P_L promoter and other promoters known to control expression of genes in prokaryotic or eukaryotic cells or their viruses. The expression vector also contains a ribosome binding site for translation initiation and a transcription terminator. The vector may also include appropriate sequences for amplifying expression.

In addition, the expression vectors preferably contain at least one selectable marker gene to provide a phenotypic trait for selection of transformed host cells. Such markers include dihydrofolate reductase (DHFR) or neomycin resistance for eukaryotic cell culture, and tetracycline or ampicillin resistance for culturing in *E. coli* and other bacteria.

The vector containing the appropriate DNA sequence as herein above described, as well as an appropriate promoter or control sequence, may be employed to transform an appropriate host to permit the host to express the protein. Representative examples of appropriate hosts, include but are not limited to: bacterial cells, such as *E. coli*, *Salmonella typhimurium*, and *Streptomyces*; fungal cells, such as yeast; insect cells, such as *Drosophila S2* and *Spodoptera Sf9*; animal cells such as CHO, COS, and Bowes melanoma; and plant cells. The selection of an appropriate host is deemed to be within the scope of those skilled in the art from the teachings herein.

More particularly, the present invention also includes recombinant constructs comprising one or more of the sequences as broadly described above. The constructs comprise a vector, such as a plasmid or viral vector, into which a sequence of the invention has been inserted, in a forward or reverse orientation. In a preferred aspect of this embodiment, the construct further comprises regulatory sequences, including, for example, a promoter, operably linked to the sequence. Large numbers of suitable vectors and promoters are known to those of skill in the art, and are commercially available. The following vectors are provided by way of example - bacterial: pQE70, pQE60 and pQE-9, available from Qiagen; pBS vectors, Phagescript vectors, Bluescript vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from Stratagene; and ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia. Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. Other suitable vectors will be readily apparent to the skilled artisan.

Promoter regions can be selected from any desired gene using CAT (chloramphenicol transferase) vectors or other vectors with selectable markers. Two appropriate vectors are pKK232-8 and pCM7. Particular named bacterial promoters include *lacI*, *lacZ*, T3, T7, gpt, lambda P_R, P_L and trp. Eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early and late SV40, LTRs from retrovirus, and mouse metallothionein-I. Selection of the appropriate vector and promoter is well within the level of ordinary skill in the art.

In a further embodiment, the present invention relates to host cells containing the above-described construct. The host cell can be a higher eukaryotic cell, such as a mammalian cell, or a lower eukaryotic cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-Dextran mediated transfection, electroporation, transduction, infection, or other methods (Davis, L., *et al.*, *Basic Methods in Molecular Biology* (1986)).

The constructs in host cells can be used in a conventional manner to produce the gene product encoded by the recombinant sequence. Alternatively, the polypeptides of the invention can be synthetically produced by conventional peptide synthesizers.

Mature proteins can be expressed in mammalian cells, yeast, bacteria, or other cells under the control of appropriate promoters. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention. Appropriate cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described by Sambrook, *et al.*, *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1989), the disclosure of which is hereby incorporated by reference.

Transcription of a DNA encoding the polypeptides of the present invention by higher eukaryotes is increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from

10 to 300 bp, that act on a promoter to increase its transcription. Examples include the SV40 enhancer on the late side of the replication origin (bp 100 to 270), a cytomegalovirus early promoter enhancer, a polyoma enhancer on the late side of the replication origin, and adenovirus enhancers.

5 Generally, recombinant expression vectors will include origins of replication and selectable markers permitting transformation of the host cell, *e.g.*, the ampicillin resistance gene of *E. coli* and *S. cerevisiae* TRP1 gene, and a promoter derived from a highly-expressed gene to direct transcription of a downstream structural sequence. Such promoters can be derived from operons encoding glycolytic enzymes such as 3-phosphoglycerate kinase (PGK), α -factor, acid phosphatase, or heat shock proteins, among others. The heterologous structural sequence is assembled in appropriate phase with translation initiation and termination sequences, and preferably, a leader sequence capable of directing secretion of translated protein into the periplasmic space or extracellular medium. Optionally, the heterologous sequence can encode a fusion protein including an N-terminal identification peptide imparting desired characteristics, *e.g.*, stabilization or simplified purification of expressed recombinant product.

10 Useful expression vectors for bacterial use are constructed by inserting a structural DNA sequence encoding a desired protein together with suitable translation initiation and termination signals in operable reading phase with a functional promoter. The vector will comprise one or more phenotypic selectable markers and an origin of replication to ensure maintenance of the vector and to, if desirable, provide amplification within the host. Suitable prokaryotic hosts for transformation include *E. coli*, *Bacillus subtilis*, *Salmonella typhimurium* and various species within the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*, although others may also be employed as a matter of choice.

15 As a representative but nonlimiting example, useful expression vectors for bacterial use can comprise a selectable marker and bacterial origin of replication derived from commercially available plasmids comprising genetic elements of the well known cloning vector pBR322 (ATCC 37017). Such

commercial vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and GEM1 (Promega Biotec, Madison, WI, USA). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed.

5 Following transformation of a suitable host strain and growth of the host strain to an appropriate cell density, the selected promoter is derepressed by appropriate means (*e.g.*, temperature shift or chemical induction) and cells are cultured for an additional period.

10 Cells are typically harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification.

15 Microbial cells employed in expression of proteins can be disrupted by any convenient method, well known to those skilled in the art, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents.

20 Various mammalian cell culture systems can also be employed to express recombinant protein. Examples of mammalian expression systems include the COS-7 lines of monkey kidney fibroblasts, described by Gluzman, *Cell* 23:175 (1981), and other cell lines capable of expressing a compatible vector, for example, the C127, 3T3, CHO, HeLa and BHK cell lines. Mammalian expression vectors will comprise an origin of replication, a suitable promoter and enhancer, and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences. DNA sequences derived from the SV40 viral genome, for example, SV40 origin, early promoter, enhancer, splice, and polyadenylation sites may be used to provide the required nontranscribed genetic elements.

25 The polypeptides can be recovered and purified from recombinant cell cultures by methods used heretofore, including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxyapatite chromatography and lectin
30 chromatography. It is preferred to have low concentrations (approximately 0.1-

5mM) of calcium ion present during purification (Price *et al.*, *J. Biol. Chem.* 244:917 (1969)). Protein refolding steps can be used, as necessary, in completing configuration of the mature protein. Finally, high performance liquid chromatography (HPLC) can be employed for final purification steps.

5 The polypeptides of the present invention may be a naturally purified product, or a product of chemical synthetic procedures, or produced by recombinant techniques from a prokaryotic or eukaryotic host (for example, by bacterial, yeast, higher plant, insect and mammalian cells in culture). Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated with mammalian or other eukaryotic carbohydrates or may be non-glycosylated. Polypeptides of the invention may also include an initial methionine amino acid residue.

10 As shown in Figures 12 and 13, the VEGF2 polypeptide of SEQ ID NO:2, minus the initial 46 amino acids, is a potent mitogen for vascular endothelial cells and stimulates their growth and proliferation. The results of a Northern blot analysis performed for the VEGF2 nucleic acid sequence encoding this polypeptide wherein 20 μ g of RNA from several human tissues were probed with ³²P-VEGF2, illustrates that this protein is actively expressed in the heart and lung which is further evidence of mitogenic activity.

15 Accordingly, VEGF2, or biologically active portions thereof, may be employed to promote angiogenesis, for example, to stimulate the growth of transplanted tissue where coronary bypass surgery is performed. VEGF2 may also be employed to promote wound healing, particularly to re-vascularize damaged tissues or stimulate collateral blood flow during ischemia and where new capillary angiogenesis is desired. VEGF2 may be employed to treat full-thickness wounds such as dermal ulcers, including pressure sores, venous ulcers, and diabetic ulcers. In addition, VEGF2 may be employed to treat full-thickness burns and injuries where a skin graft or flap is used to repair such burns and injuries. VEGF2 may also be employed for use in plastic surgery, for example, for the repair of lacerations, burns, or other trauma.

20

25

30

Along these same lines, VEGF2 may also be employed to induce the growth of damaged bone, periodontium or ligament tissue. VEGF2 may also be employed for regenerating supporting tissues of the teeth, including cementum and periodontal ligament, that have been damaged by, *e.g.*, periodental disease or trauma.

Since angiogenesis is important in keeping wounds clean and non-infected, VEGF2 may be employed in association with surgery and following the repair of cuts. It may also be employed for the treatment of abdominal wounds where there is a high risk of infection.

VEGF2 may be employed for the promotion of endothelialization in vascular graft surgery. In the case of vascular grafts using either transplanted or synthetic material, VEGF2 can be applied to the surface of the graft or at the junction to promote the growth of vascular endothelial cells. VEGF2 may also be employed to repair damage of myocardial tissue as a result of myocardial infarction. VEGF2 may also be employed to repair the cardiac vascular system after ischemia. VEGF2 may also be employed to treat damaged vascular tissue as a result of coronary artery disease and peripheral and CNS vascular disease.

VEGF2 may also be employed to coat artificial prostheses or natural organs which are to be transplanted in the body to minimize rejection of the transplanted material and to stimulate vascularization of the transplanted materials.

VEGF2 may also be employed for vascular tissue repair, for example, that occurring during arteriosclerosis and required following balloon angioplasty where vascular tissues are damaged.

VEGF2 nucleic acid sequences and VEGF2 polypeptides may also be employed for *in vitro* purposes related to scientific research, synthesis of DNA and manufacture of DNA vectors, and for the production of diagnostics and therapeutics to treat human disease. For example, VEGF2 may be employed for *in vitro* culturing of vascular endothelial cells, where it is added to the conditional medium in a concentration from 10 pg/ml to 10 ng/ml.

Fragments of the full length VEGF2 gene may be used as a hybridization probe for a cDNA library to isolate other genes which have a high sequence similarity to the gene or similar biological activity. Probes of this type generally have at least 50 base pairs, although they may have a greater number of bases. The probe may also be used to identify a cDNA clone corresponding to a full length transcript and a genomic clone or clones that contain the complete VEGF2 gene including regulatory and promotor regions, exons, and introns. An example of a screen comprises isolating the coding region of the VEGF2 gene by using the known DNA sequence to synthesize an oligonucleotide probe. Labeled oligonucleotides having a sequence complementary to that of the gene of the present invention are used to screen a library of human cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes to.

This invention provides methods for identification of VEGF2 receptors. The gene encoding the receptor can be identified by numerous methods known to those of skill in the art, for example, ligand panning and FACS sorting (Coligan *et al.*, *Current Protocols in Immun.*, 1(2), Chapter 5, (1991)). Preferably, expression cloning is employed wherein polyadenylated RNA is prepared from a cell responsive to VEGF2, and a cDNA library created from this RNA is divided into pools and used to transfect COS cells or other cells that are not responsive to VEGF2. Transfected cells which are grown on glass slides are exposed to labeled VEGF2. VEGF2 can be labeled by a variety of means including iodination or inclusion of a recognition site for a site-specific protein kinase. Following fixation and incubation, the slides are subjected to autoradiographic analysis. Positive pools are identified and sub-pools are prepared and retransfected using an iterative sub-pooling and rescreening process, eventually yielding a single clone that encodes the putative receptor.

As an alternative approach for receptor identification, labeled VEGF2 can be photoaffinity linked with cell membrane or extract preparations that express the receptor molecule. Cross-linked material is resolved by PAGE and exposed to X-ray film. The labeled complex containing VEGF2 is then excised, resolved

into peptide fragments, and subjected to protein microsequencing. The amino acid sequence obtained from microsequencing would be used to design a set of degenerate oligonucleotide probes to screen a cDNA library to identify the gene encoding the putative receptor.

5 This invention is also related to a method of screening compounds to identify those which are VEGF2 agonists or antagonists. An example of such a method takes advantage of the ability of VEGF2 to significantly stimulate the proliferation of human endothelial cells in the presence of the comitogen Con A. Endothelial cells are obtained and cultured in 96-well flat-bottomed culture plates (Costar, Cambridge, MA) in a reaction mixture supplemented with Con-A (Calbiochem, La Jolla, CA). Con-A, polypeptides of the present invention and the compound to be screened are added. After incubation at 37°C, cultures are pulsed with 1 μ Ci of 3 [H]thymidine (5 Ci/mmol; 1 Ci = 37 BGq; NEN) for a sufficient time to incorporate the 3 [H] and harvested onto glass fiber filters (Cambridge Technology, Watertown, MA). Mean 3 [H]-thymidine incorporation (cpm) of triplicate cultures is determined using a liquid scintillation counter (Beckman Instruments, Irvine, CA). Significant 3 [H]thymidine incorporation, as compared to a control assay where the compound is excluded, indicates stimulation of endothelial cell proliferation.

10
15
20 To assay for antagonists, the assay described above is performed and the ability of the compound to inhibit 3 [H]thymidine incorporation in the presence of VEGF2 indicates that the compound is an antagonist to VEGF2. Alternatively, VEGF2 antagonists may be detected by combining VEGF2 and a potential antagonist with membrane-bound VEGF2 receptors or recombinant receptors under appropriate conditions for a competitive inhibition assay. VEGF2 can be labeled, such as by radioactivity, such that the number of VEGF2 molecules bound to the receptor can determine the effectiveness of the potential antagonist.

25
30 Alternatively, the response of a known second messenger system following interaction of VEGF2 and receptor would be measured and compared in the presence or absence of the compound. Such second messenger systems

include but are not limited to, cAMP guanylate cyclase, ion channels or phosphoinositide hydrolysis. In another method, a mammalian cell or membrane preparation expressing the VEGF2 receptor is incubated with labeled VEGF2 in the presence of the compound. The ability of the compound to enhance or block this interaction could then be measured.

Potential VEGF2 antagonists include an antibody, or in some cases, an oligonucleotide, which bind to the polypeptide and effectively eliminate VEGF2 function. Alternatively, a potential antagonist may be a closely related protein which binds to VEGF2 receptors, however, they are inactive forms of the polypeptide and thereby prevent the action of VEGF2. Examples of these antagonists include a negative dominant mutant of the VEGF2 polypeptide, for example, one chain of the hetero-dimeric form of VEGF2 may be dominant and may be mutated such that biological activity is not retained. An example of a negative dominant mutant includes truncated versions of a dimeric VEGF2 which is capable of interacting with another dimer to form wild type VEGF2, however, the resulting homo-dimer is inactive and fails to exhibit characteristic VEGF activity.

Another potential VEGF2 antagonist is an antisense construct prepared using antisense technology. Antisense technology can be used to control gene expression through triple-helix formation or antisense DNA or RNA, both of which methods are based on binding of a polynucleotide to DNA or RNA. For example, the 5' coding portion of the polynucleotide sequence, which encodes for the mature polypeptides of the present invention, is used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription (triple helix -see Lee *et al.*, *Nucl. Acids Res.*6:3073 (1979); Cooney *et al.*, *Science* 241:456 (1988); and Dervan *et al.*, *Science* 251:1360 (1991)), thereby preventing transcription and the production of VEGF2. The antisense RNA oligonucleotide hybridizes to the mRNA *in vivo* and blocks translation of the mRNA molecule into the VEGF2 polypeptide (Antisense -

Okano, *J. Neurochem.*56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988)). The oligonucleotides described above can also be delivered to cells such that the antisense RNA or DNA may be expressed *in vivo* to inhibit production of VEGF2.

Potential VEGF2 antagonists also include small molecules which bind to and occupy the active site of the polypeptide thereby making the catalytic site inaccessible to substrate such that normal biological activity is prevented. Examples of small molecules include but are not limited to small peptides or peptide-like molecules.

The antagonists may be employed to limit angiogenesis necessary for solid tumor metastasis. The identification of VEGF2 can be used for the generation of certain inhibitors of vascular endothelial growth factor. Since angiogenesis and neovascularization are essential steps in solid tumor growth, inhibition of angiogenic activity of the vascular endothelial growth factor is very useful to prevent the further growth, retard, or even regress solid tumors. Although the level of expression of VEGF2 is extremely low in normal tissues including breast, it can be found expressed at moderate levels in at least two breast tumor cell lines that are derived from malignant tumors. It is, therefore, possible that VEGF2 is involved in tumor angiogenesis and growth.

Gliomas are also a type of neoplasia which may be treated with the antagonists of the present invention.

The antagonists may also be used to treat chronic inflammation caused by increased vascular permeability. In addition to these disorders, the antagonists may also be employed to treat retinopathy associated with diabetes, rheumatoid arthritis and psoriasis.

The antagonists may be employed in a composition with a pharmaceutically acceptable carrier, *e.g.*, as hereinafter described.

The VEGF2 polypeptides and agonists and antagonists may be employed in combination with a suitable pharmaceutical carrier. Such compositions

comprise a therapeutically effective amount of the polypeptide or agonist or antagonist, and a pharmaceutically acceptable carrier or excipient. Such a carrier includes but is not limited to saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The formulation should suit the mode of administration.

The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. In addition, the pharmaceutical compositions may be employed in conjunction with other therapeutic compounds.

The pharmaceutical compositions may be administered in a convenient manner such as by the topical, intravenous, intraperitoneal, intramuscular, intratumor, subcutaneous, intranasal or intradermal routes. The pharmaceutical compositions are administered in an amount which is effective for treating and/or prophylaxis of the specific indication. In general, the pharmaceutical compositions are administered in an amount of at least about 10 $\mu\text{g/kg}$ body weight and in most cases they will be administered in an amount not in excess of about 8 mg/Kg body weight per day. In most cases, the dosage is from about 10 $\mu\text{g/kg}$ to about 1 mg/kg body weight daily, taking into account the routes of administration, symptoms, etc.

The VEGF2 polypeptides, and agonists or antagonists which are polypeptides may also be employed in accordance with the present invention by expression of such polypeptide *in vivo*, which is often referred to as "gene therapy."

Thus, for example, cells such as bone marrow cells may be engineered with a polynucleotide (DNA or RNA) encoding for the polypeptide *ex vivo*, the engineered cells are then provided to a patient to be treated with the polypeptide.

Such methods are well-known in the art. For example, cells may be engineered by procedures known in the art by use of a retroviral particle containing RNA encoding the polypeptide of the present invention.

Similarly, cells may be engineered *in vivo* for expression of a polypeptide *in vivo*, for example, by procedures known in the art. As known in the art, a producer cell for producing a retroviral particle containing RNA encoding a polypeptide of the present invention may be administered to a patient for engineering cells *in vivo* and expression of the polypeptide *in vivo*. These and other methods for administering a polypeptide of the present invention by such methods should be apparent to those skilled in the art from the teachings of the present invention. For example, the expression vehicle for engineering cells may be other than a retroviral particle, for example, an adenovirus, which may be used to engineer cells *in vivo* after combination with a suitable delivery vehicle.

Retroviruses from which the retroviral plasmid vectors hereinabove mentioned may be derived include, but are not limited to, Moloney Murine Leukemia Virus, spleen necrosis virus, retroviruses such as Rous Sarcoma Virus, Harvey Sarcoma Virus, avian leukosis virus, gibbon ape leukemia virus, human immunodeficiency virus, adenovirus, Myeloproliferative Sarcoma Virus, and mammary tumor virus. In one embodiment, the retroviral plasmid vector is derived from Moloney Murine Leukemia Virus.

The vector includes one or more promoters. Suitable promoters which may be employed include, but are not limited to, the retroviral LTR; the SV40 promoter; and the human cytomegalovirus (CMV) promoter described in Miller *et al.*, *Biotechniques* 7:980-990 (1989), or any other promoter (*e.g.*, cellular promoters such as eukaryotic cellular promoters including, but not limited to, the histone, pol III, and β -actin promoters). Other viral promoters which may be employed include, but are not limited to, adenovirus promoters, thymidine kinase (TK) promoters, and B19 parvovirus promoters. The selection of a suitable promoter will be apparent to those skilled in the art from the teachings contained herein.

5 The nucleic acid sequence encoding the polypeptide of the present invention is under the control of a suitable promoter. Suitable promoters which may be employed include, but are not limited to, adenoviral promoters, such as the adenoviral major late promoter; or heterologous promoters, such as the cytomegalovirus (CMV) promoter; the respiratory syncytial virus (RSV) promoter; inducible promoters, such as the MMT promoter, the metallothionein promoter; heat shock promoters; the albumin promoter; the ApoAI promoter; human globin promoters; viral thymidine kinase promoters, such as the Herpes Simplex thymidine kinase promoter; retroviral LTRs (including the modified retroviral LTRs hereinabove described); the β -actin promoter; and human growth hormone promoters. The promoter also may be the native promoter which controls the gene encoding the polypeptide.

10 The retroviral plasmid vector is employed to transduce packaging cell lines to form producer cell lines. Examples of packaging cells which may be transfected include, but are not limited to, the PE501, PA317, ψ -2, ψ -AM, PA12, T19-14X, VT-19-17-H2, ψ CRE, ψ CRIP, GP+E-86, GP+envAm12, and DAN cell lines as described in Miller, *Human Gene Therapy* 1:5-14 (1990), which is incorporated herein by reference in its entirety. The vector may transduce the packaging cells through any means known in the art. Such means include, but are not limited to, electroporation, the use of liposomes, and CaPO_4 precipitation. In one alternative, the retroviral plasmid vector may be encapsulated into a liposome, or coupled to a lipid, and then administered to a host.

20 The producer cell line generates infectious retroviral vector particles which include the nucleic acid sequence(s) encoding the polypeptides. Such retroviral vector particles then may be employed, to transduce eukaryotic cells, either *in vitro* or *in vivo*. The transduced eukaryotic cells will express the nucleic acid sequence(s) encoding the polypeptide. Eukaryotic cells which may be transduced include, but are not limited to, embryonic stem cells, embryonic carcinoma cells, as well as hematopoietic stem cells, hepatocytes, fibroblasts, myoblasts, keratinocytes, endothelial cells, and bronchial epithelial cells.

This invention is also related to the use of the VEGF2 gene as part of a diagnostic assay for detecting diseases or susceptibility to diseases related to the presence of mutations in VEGF2 nucleic acid sequences.

5 Individuals carrying mutations in the VEGF2 gene may be detected at the DNA level by a variety of techniques. Nucleic acids for diagnosis may be obtained from a patient's cells, such as from blood, urine, saliva, tissue biopsy and autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR (Saiki *et al.*, *Nature* 324:163-166 (1986)) prior to analysis. RNA or cDNA may also be used for the same purpose. As an example, PCR primers complementary to the nucleic acid encoding VEGF2 can be used to identify and analyze VEGF2 mutations. For example, deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to radiolabeled VEGF2 RNA or alternatively, radiolabeled VEGF2 antisense DNA sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase A digestion or by differences in melting temperatures.

10
15
20 Genetic testing based on DNA sequence differences may be achieved by detection of alteration in electrophoretic mobility of DNA fragments in gels with or without denaturing agents. Small sequence deletions and insertions can be visualized by high resolution gel electrophoresis. DNA fragments of different sequences may be distinguished on denaturing formamide gradient gels in which the mobilities of different DNA fragments are retarded in the gel at different positions according to their specific melting or partial melting temperatures (see, *e.g.*, Myers *et al.*, *Science* 230:1242 (1985)).

25 Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method (*e.g.*, Cotton *et al.*, *PNAS, USA* 85:4397-4401 (1985)).

30 Thus, the detection of a specific DNA sequence may be achieved by methods such as hybridization, RNase protection, chemical cleavage, direct DNA

sequencing or the use of restriction enzymes, (*e.g.*, Restriction Fragment Length Polymorphisms (RFLP)) and Southern blotting of genomic DNA.

In addition to more conventional gel-electrophoresis and DNA sequencing, mutations can also be detected by *in situ* analysis.

5 The present invention also relates to a diagnostic assay for detecting altered levels of VEGF2 protein in various tissues since an over-expression of the proteins compared to normal control tissue samples may detect the presence of a disease or susceptibility to a disease, for example, abnormal cellular differentiation. Assays used to detect levels of VEGF2 protein in a sample
10 derived from a host are well-known to those of skill in the art and include radioimmunoassays, competitive-binding assays, Western Blot analysis, ELISA assays and "sandwich" assay. An ELISA assay (Coligan *et al.*, *Current Protocols in Immunology* 1(2), Chapter 6, (1991)) initially comprises preparing an antibody specific to the VEGF2 antigen, preferably a monoclonal antibody. In addition a
15 reporter antibody is prepared against the monoclonal antibody. To the reporter antibody is attached a detectable reagent such as radioactivity, fluorescence or, in this example, a horseradish peroxidase enzyme. A sample is removed from a host and incubated on a solid support, *e.g.* a polystyrene dish, that binds the proteins in the sample. Any free protein binding sites on the dish are then
20 covered by incubating with a non-specific protein, such as, bovine serum albumen. Next, the monoclonal antibody is incubated in the dish during which time the monoclonal antibodies attach to any VEGF2 proteins attached to the polystyrene dish. All unbound monoclonal antibody is washed out with buffer. The reporter antibody linked to horseradish peroxidase is placed in the dish
25 resulting in binding of the reporter antibody to any monoclonal antibody bound to VEGF2. Unattached reporter antibody is then washed out. Peroxidase substrates are then added to the dish and the amount of color developed in a given time period is a measurement of the amount of VEGF2 protein present in a given volume of patient sample when compared against a standard curve.

A competition assay may be employed wherein antibodies specific to VEGF2 are attached to a solid support. Polypeptides of the present invention are then labeled, for example, by radioactivity, and a sample derived from the host are passed over the solid support and the amount of label detected, for example by liquid scintillation chromatography, can be correlated to a quantity of VEGF2 in the sample.

A "sandwich" assay is similar to an ELISA assay. In a "sandwich" assay VEGF2 is passed over a solid support and binds to antibody attached to a solid support. A second antibody is then bound to the VEGF2. A third antibody which is labeled and specific to the second antibody is then passed over the solid support and binds to the second antibody and an amount can then be quantified.

The sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. Moreover, there is a current need for identifying particular sites on the chromosome. Few chromosome marking reagents based on actual sequence data (repeat polymorphism's) are presently available for marking chromosomal location. The mapping of DNAs to chromosomes according to the present invention is an important first step in correlating those sequences with genes associated with disease.

Briefly, sequences can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp) from the cDNA. Computer analysis of the cDNA is used to rapidly select primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers are then used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the primer will yield an amplified fragment.

PCR mapping of somatic cell hybrids is a rapid procedure for assigning a particular DNA to a particular chromosome. Using the present invention with the same oligonucleotide primers, sublocalization can be achieved with panels of fragments from specific chromosomes or pools of large genomic clones in an

analogous manner. Other mapping strategies that can similarly be used to map to its chromosome include *in situ* hybridization, prescreening with labeled flow-sorted chromosomes and preselection by hybridization to construct chromosome specific-cDNA libraries.

5 Fluorescence *in situ* hybridization (FISH) of a cDNA clone to a metaphase chromosomal spread can be used to provide a precise chromosomal location in one step. This technique can be used with probes from the cDNA as short as 50 or 60 base pairs. For a review of this technique, see Verma *et al.*, *Human Chromosomes: a Manual of Basic Techniques*, Pergamon Press, New York (1988).

10 Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes).

15 Next, it is necessary to determine the differences in the cDNA or genomic sequence between affected and unaffected individuals. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

20 With current resolution of physical mapping and genetic mapping techniques, a cDNA precisely localized to a chromosomal region associated with the disease could be one of between 50 and 500 potential causative genes. (This assumes 1 megabase mapping resolution and one gene per 20 kb).

25 Comparison of affected and unaffected individuals generally involves first looking for structural alterations in the chromosomes, such as deletions or translocations that are visible from chromosome spreads or detectable using PCR based on that cDNA sequence. Ultimately, complete sequencing of genes from

several individuals is required to confirm the presence of a mutation and to distinguish mutations from polymorphisms.

5 The present invention is further directed to inhibiting VEGF2 *in vivo* by the use of antisense technology. Antisense technology can be used to control gene expression through triple-helix formation or antisense DNA or RNA, both of which methods are based on binding of a polynucleotide to DNA or RNA. For example, the 5' coding portion of the mature polynucleotide sequence, which encodes for the polypeptide of the present invention, is used to design an antisense RNA oligonucleotide of from 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription (triple helix -- see Lee *et al.*, *Nucl. Acids Res.* 6:3073 (1979); Cooney *et al.*, *Science* 241:456 (1988); and Dervan *et al.* *Science*, 251:1360 (1991), thereby preventing transcription and the production of VEGF2. The antisense RNA oligonucleotide hybridizes to the mRNA *in vivo* and blocks translation of an mRNA molecule into the VEGF2 (antisense -- Okano, *J. Neurochem.* 56:560 (1991); *Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression*, CRC Press, Boca Raton, FL (1988)).

10
15
20 Alternatively, the oligonucleotides described above can be delivered to cells by procedures in the art such that the anti-sense RNA or DNA may be expressed *in vivo* to inhibit production of VEGF2 in the manner described above.

25 Antisense constructs to VEGF2, therefore, may inhibit the angiogenic activity of the VEGF2 and prevent the further growth or even regress solid tumors, since angiogenesis and neovascularization are essential steps in solid tumor growth. These antisense constructs may also be used to treat rheumatoid arthritis, psoriasis and diabetic retinopathy which are all characterized by abnormal angiogenesis.

30 The polypeptides, their fragments or other derivatives, or analogs thereof, or cells expressing them can be used as an immunogen to produce antibodies thereto. These antibodies can be, for example, polyclonal or monoclonal antibodies. The present invention also includes chimeric, single chain, and

humanized antibodies, as well as Fab fragments, or the product of an Fab expression library. Various procedures known in the art may be used for the production of such antibodies and fragments.

Antibodies generated against the polypeptide corresponding to a sequence of the present invention can be obtained by direct injection of the polypeptide into an animal or by administering the polypeptide to an animal, preferably a nonhuman. The antibody so obtained will then bind the polypeptide itself. In this manner, even a sequence encoding only a fragment of the polypeptide can be used to generate antibodies binding the whole native polypeptide. Such antibodies can then be used to isolate the polypeptide from tissue expressing that polypeptide.

For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler and Milstein, *Nature* 256:495-497 (1975)), the trioma technique, the human B-cell hybridoma technique (Kozbor *et al.*, *Immunology Today* 4:72 (1983)), and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole *et al.*, in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc. (1985), pp. 77-96).

Techniques described for the production of single chain antibodies (U.S. Patent 4,946,778) can be adapted to produce single chain antibodies to immunogenic polypeptide products of this invention.

Neutralization antibodies can be identified and applied to mask the vascular endothelial growth factor, and that has been shown in mice model systems against VEGF. VEGF2 can also be inactivated by certain dominant negative mutants within the gene itself. It is known that both PDGF α and β form either heterodimers or homodimers, and VEGF forms homodimers. Similar interaction between VEGF2 could be expected. These antibodies therefore may be used to block the angiogenic activity of VEGF2 and retard the growth of solid tumors. These antibodies may also be used to treat inflammation caused by the increased vascular permeability which results from the presence of VEGF2.

These antibodies may further be used in an immunoassay to detect the presence of tumors in certain individuals. Enzyme immunoassay can be performed from the blood sample of an individual. Elevated levels of VEGF2 can be considered diagnostic of cancer.

5 The present invention is also directed to antagonist/inhibitors of the polypeptides of the present invention. The antagonist/inhibitors are those which inhibit or eliminate the function of the polypeptide.

10 Thus, for example, antagonists bind to a polypeptide of the present invention and inhibit or eliminate its function. The antagonist, for example, could be an antibody against the polypeptide which binds to the polypeptide or, in some cases, an oligonucleotide. An example of an inhibitor is a small molecule which binds to and occupies the catalytic site of the polypeptide thereby making the catalytic site inaccessible to substrate such that normal biological activity is prevented. Examples of small molecules include but are not limited to small peptides or peptide-like molecules.

15 Truncated versions of VEGF2 can also be produced that are capable of interacting with wild type VEGF2 to form dimers that fail to activate endothelial cell growth, therefore inactivating the endogenous VEGF2. Or, mutant forms of VEGF2 form dimers themselves and occupy the ligand binding domain of the proper tyrosine kinase receptors on the target cell surface, but fail to activate cell growth.

20 Alternatively, antagonists to the polypeptides of the present invention may be employed which bind to the receptors to which a polypeptide of the present invention normally binds. The antagonists may be closely related proteins such that they recognize and bind to the receptor sites of the natural protein, however, they are inactive forms of the natural protein and thereby prevent the action of VEGF2 since receptor sites are occupied. In these ways, the action of the VEGF2 is prevented and the antagonist/inhibitors may be used therapeutically as an anti-tumor drug by occupying the receptor sites of tumors which are recognized by VEGF2 or by inactivating VEGF2 itself. The antagonist/inhibitors may also be

25

30

used to prevent inflammation due to the increased vascular permeability action of VEGF2. The antagonist/inhibitors may also be used to treat solid tumor growth, diabetic retinopathy, psoriasis and rheumatoid arthritis.

The antagonist/inhibitors may be employed in a composition with a pharmaceutically acceptable carrier, *e.g.*, as hereinabove described.

The present invention will be further described with reference to the following examples; however, it is to be understood that the present invention is not limited to such examples. All parts or amounts, unless otherwise specified, are by weight.

In order to facilitate understanding of the following examples, certain frequently occurring methods and/or terms will be described.

"Plasmids" are designated by a lower case p preceded and/or followed by capital letters and/or numbers. The starting plasmids herein are either commercially available, publicly available on an unrestricted basis, or can be constructed from available plasmids in accord with published procedures. In addition, equivalent plasmids to those described are known in the art and will be apparent to the ordinarily skilled artisan.

"Digestion" of DNA refers to catalytic cleavage of the DNA with a restriction enzyme that acts only at certain sequences in the DNA. The various restriction enzymes used herein are commercially available and their reaction conditions, cofactors and other requirements were used as would be known to the ordinarily skilled artisan. For analytical purposes, typically 1 μ g of plasmid or DNA fragment is used with about 2 units of enzyme in about 20 μ l of buffer solution. For the purpose of isolating DNA fragments for plasmid construction, typically 5 to 50 μ g of DNA are digested with 20 to 250 units of enzyme in a larger volume. Appropriate buffers and substrate amounts for particular restriction enzymes are specified by the manufacturer. Incubation times of about 1 hour at 37°C are ordinarily used, but may vary in accordance with the supplier's instructions. After digestion the reaction is electrophoresed directly on a polyacrylamide gel to isolate the desired fragment.

Size separation of the cleaved fragments is performed using 8 percent polyacrylamide gel described by Goeddel, D. *et al.*, *Nucleic Acids Res.* 8:4057 (1980).

"Oligonucleotides" refer to either a single stranded polydeoxynucleotide or two complementary polydeoxynucleotide strands, which may be chemically synthesized. Such synthetic oligonucleotides have no 5' phosphate and thus will not ligate to another oligonucleotide without adding a phosphate with an ATP in the presence of a kinase. A synthetic oligonucleotide will ligate to a fragment that has not been dephosphorylated.

"Ligation" refers to the process of forming phosphodiester bonds between two double stranded nucleic acid fragments (Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1989), p. 146). Unless otherwise provided, ligation may be accomplished using known buffers and conditions with 10 units of T4 DNA ligase ("ligase") per 0.5 μ g of approximately equimolar amounts of the DNA fragments to be ligated.

Unless otherwise stated, transformation was performed as described by the method of Graham, F. and Van der Eb, A., *Virology* 52:456-457 (1973).

Example 1

Expression Pattern of VEGF2 in Human Tissues and Breast Cancer Cell Lines

Northern blot analysis was carried out to examine the levels of expression of VEGF2 in human tissues and breast cancer cell lines in human tissues. Total cellular RNA samples were isolated with RNeasyTM B system (Qiagen Laboratories, Inc.). About 10 μ g of total RNA isolated from each breast tissue and cell line specified was separated on 1% agarose gel and blotted onto a nylon filter, (Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1989)). The labeling reaction was done according to the Stratagene Prime-It kit with 50 ng

DNA fragment. The labeled DNA was purified with a Select-G-50 column from 5 Prime → 3 Prime, Inc (Boulder, CO). The filter was then hybridized with a radioactive labeled full length VEGF2 gene at 1,000,000 cpm/ml in 0.5 M NaPO₄ and 7 % SDS overnight at 65°C. After washing twice at room temperature and twice at 60°C with 0.5 X SSC, 0.1 % SDS, the filters were then exposed at -70°C overnight with an intensifying screen. A message of 1.6 Kd was observed in 2 breast cancer cell lines. FIG. 5, lane #4 represents a very tumorigenic cell line that is estrogen independent for growth.

Also, 10 µg of total RNA from 10 human adult tissues were separated on an agarose gel and blotted onto a nylon filter. The filter was then hybridized with radioactively labeled VEGF2 probe in 7% SDS, 0.5 M NaPO₄, pH 7.2; 1% BSA overnight at 65°C. Following washing in 0.2 X SSC at 65°C, the filter was exposed to film for 24 days at -70°C with intensifying screen. See FIG. 6.

Example 2

Expression of the Truncated Form of VEGF2 (SEQ ID NO:4) by in vitro Transcription and Translation

The VEGF2 cDNA was transcribed and translated *in vitro* to determine the size of the translatable polypeptide encoded by the truncated form of VEGF2 and a partial VEGF2 cDNA. The two inserts of VEGF2 in the pBluescript SK vector were amplified by PCR with three pairs of primers, 1) M13-reverse and forward primers; 2) M13-reverse primer and VEGF primer F4; and 3) M13-reverse primer and VEGF primer F5. The sequence of these primers are as follows.

M13-2 reverse primer:

5'-ATGCTTCCGGCTCGTATG-3' (SEQ ID NO: 9)

This sequence is located upstream of the 5' end of the VEGF2 cDNA insert in the pBluescript vector and is in an anti-sense orientation as the cDNA. A T3 promoter sequence is located between this primer and the VEGF2 cDNA.

M13-2 forward primer:

5'GGGTTTTCCCAGTCACGAC-3' (SEQ ID NO: 10)

This sequence is located downstream of the 3' end of the VEGF2 cDNA insert in the pBluescript vector and is in an anti-sense orientation as the cDNA insert.

VEGF primer F4:

5'-CCACATGGTTCAGGAAAGACA-3' (SEQ ID NO: 11)

This sequence is located within the VEGF2 cDNA in an anti-sense orientation from bp 1259-1239, which is about 169 bp away from the 3' end of the stop codon and about 266 bp before the last nucleotide of the cDNA.

PCR reaction with all three pairs of primers produce amplified products with T3 promoter sequence in front of the cDNA insert. The first and third pairs of primers produce PCR products that encode the polypeptide of VEGF2 shown in SEQ ID NO:4. The second pair of primers produce PCR product that misses 36 amino acids coding sequence at the C-terminus of the VEGF2 polypeptide.

Approximately 0.5 μ g of PCR product from first pair of primers, 1 μ g from second pair of primers, 1 μ g from third pair of primers were used for *in vitro* transcription/translation. The *in vitro* transcription/translation reaction was performed in a 25 μ l of volume, using the T_NT™ Coupled Reticulocyte Lysate Systems (Promega, CAT# L4950). Specifically, the reaction contains 12.5 μ l of T_NT rabbit reticulocyte lysate 2 μ l of T_NT reaction buffer, 1 μ l of T3 polymerase, 1 μ l of 1 mM amino acid mixtrue (minus methionine), 4 μ l of ³⁵S-methionine (>1000 Ci/mmol, 10 mCi/ml), 1 μ l of 40 U/ul; RNasin ribonuclease inhibitor, 0.5 or 1 μ g of PCR products. Nuclease-free H₂O was added to bring the volume to 25 μ l. The reaction was incubated at 30°C for 2 hours. Five microliters of the reaction product was analyzed on a 4-20% gradient SDS-PAGE gel. After fixing in 25% isopropanol and 10% acetic acid, the gel was dried and exposed to an X-ray film overnight at 70°C.

As shown in FIG. 7, PCR products containing the truncated VEGF2 cDNA (*i.e.*, as depicted in SEQ ID NO:3) and the cDNA missing 266 bp in the 3' un-translated region (3'-UTR) produced the same length of translated products,

whose molecular weights are estimated to be 38-40 dk (lanes 1 and 3). The cDNA missing all the 3'UTR and missing sequence encoding the C-terminal 36 amino acids was translated into a polypeptide with an estimated molecular weight of 36-38 kd (lane 2).

Example 3

Cloning and Expression of VEGF2 Using the Baculovirus Expression System

The DNA sequence encoding the VEGF2 protein without 46 amino acids at the N-terminus, see ATCC No. 97149, was amplified using PCR oligonucleotide primers corresponding to the 5' and 3' sequences of the gene:

The 5' primer has the sequence TGT AAT ACG ACT CAC TAT AGG **GAT CCC** GCC ATG GAG GCC ACG GCT TAT GC (SEQ ID NO:12) and contains a BamH1 restriction enzyme site (in bold) and 17 nucleotide sequence complementary to the 5' sequence of VEGF2 (nt. 150-166).

The 3' primer has the sequence GATC **TCT AGA** TTA GCT CAT TTG TGG TCT (SEQ ID NO:13) and contains the cleavage site for the restriction enzyme XbaI and 18 nucleotides complementary to the 3' sequence of VEGF2, including the stop codon and 15 nt sequence before stop codon.

The amplified sequences were isolated from a 1% agarose gel using a commercially available kit ("GeneClean," BIO 101, Inc., La Jolla, CA). The fragment was then digested with the endonuclease BamH1 and XbaI and then purified again on a 1% agarose gel. This fragment was ligated to pAcGP67A baculovirus transfer vector (PharMingen) at the BamH1 and XbaI sites. Through this ligation, VEGF2 cDNA was cloned in frame with the signal sequence of baculovirus gp67 gene and was located at the 3' end of the signal sequence in the vector. This is designated pAcGP67A-VEGF2.

To clone VEGF2 with the signal sequence of gp67 gene to the pRG1 vector for expression, VEGF2 with the signal sequence and some upstream sequence were excised from the pAcGP67A-VEGF2 plasmid at the Xho

restriction endonuclease site located upstream of the VEGF2 cDNA and at the XbaI restriction endonuclease site by XhoI and XbaI restriction enzyme. This fragment was separated from the rest of vector on a 1% agarose gel and was purified using "GeneClean" kit. It was designated F2.

5 The PRG1 vector (modification of pVL941 vector) is used for the expression of the VEGF2 protein using the baculovirus expression system (for review see: Summers, M.D. and Smith, G.E., *"A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures,"* Texas Agricultural Experimental Station Bulletin No. 1555, (1987)). This expression vector contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus (AcMNPV) followed by the recognition sites for the restriction endonucleases BamHI, SmaI, XbaI, BglII and Asp718. A site for restriction endonuclease XhoI is located upstream of BamHI site. The sequence between XhoI and BamHI is the same as that in pAcGp67A (static on tape) vector. The polyadenylation site of the simian virus (SV)40 is used for efficient polyadenylation. For an easy selection of recombinant virus the beta-galactosidase gene from E.coli is inserted in the same orientation as the polyhedrin promoter followed by the polyadenylation signal of the polyhedrin gene. The polyhedrin sequences are flanked at both sides by viral sequences for
10
15
20 the cell-mediated homologous recombination of cotransfected wild-type viral DNA. Many other baculovirus vectors could be used in place of pRG1 such as pAc373, pVL941 and pAcIM1 (Luckow, V.A. and Summers, M.D., *Virology* 170:31-39 (1989).

25 The plasmid was digested with the restriction enzymes XhoI and XbaI and then dephosphorylated using calf intestinal phosphatase by procedures known in the art. The DNA was then isolated from a 1% agarose gel using the commercially available kit ("GeneClean" BIO 101 Inc., La Jolla, Ca.). This vector DNA is designated V2.

30 Fragment F2 and the dephosphorylated plasmid V2 were ligated with T4 DNA ligase. *E.coli* HB101 cells were then transformed and bacteria identified

that contained the plasmid (pBac gp67-VEGF2) with the VEGF2 gene using the enzymes BamHI and XbaI. The sequence of the cloned fragment was confirmed by DNA sequencing.

5 5 µg of the plasmid pBac gp67-VEGF2 was cotransfected with 1.0 µg of a commercially available linearized baculovirus ("BaculoGold™ baculovirus DNA", Pharmingen, San Diego, CA.) using the lipofectin method (Felgner et al. Proc. Natl. Acad. Sci. USA, 84:7413-7417 (1987)).

10 1 µg of BaculoGold™ virus DNA and 5 µg of the plasmid pBac gp67-VEGF2 were mixed in a sterile well of a microtiter plate containing 50 µl of serum free Grace's medium (Life Technologies Inc., Gaithersburg, MD). Afterwards 10 µl Lipofectin plus 90 µl Grace's medium were added, mixed and incubated for 15 minutes at room temperature. Then the transfection mixture was added dropwise to the Sf9 insect cells (ATCC CRL 1711) seeded in a 35 mm tissue culture plate with 1 ml Grace's medium without serum. The plate was
15 rocked back and forth to mix the newly added solution. The plate was then incubated for 5 hours at 27°C. After 5 hours the transfection solution was removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum was added. The plate was put back into an incubator and cultivation continued at 27°C for four days.

20 After four days the supernatant was collected and a plaque assay performed similar as described by Summers and Smith, *supra*. As a modification an agarose gel with "Blue Gal" (Life Technologies Inc., Gaithersburg) was used which allows an easy isolation of blue stained plaques. (A detailed description of a "plaque assay" can also be found in the user's guide for insect cell culture and
25 baculovirology distributed by Life Technologies Inc., Gaithersburg, page 9-10).

30 Four days after the serial dilution, the virus was added to the cells, blue stained plaques were picked with the tip of an Eppendorf pipette. The agar containing the recombinant viruses was then resuspended in an Eppendorf tube containing 200 µl of Grace's medium. The agar was removed by a brief centrifugation and the supernatant containing the recombinant baculovirus was

used to infect Sf9 cells seeded in 35 mm dishes. Four days later the supernatants of these culture dishes were harvested and then stored at 4°C.

Sf9 cells were grown in Grace's medium supplemented with 10% heat-inactivated FBS. The cells were infected with the recombinant baculovirus V-gp67-VEGF2 at a multiplicity of infection (MOI) of 1. Six hours later the medium was removed and replaced with SF900 II medium minus methionine and cysteine (Life Technologies Inc., Gaithersburg). 42 hours later 5 µCi of ³⁵S-methionine and 5 µCi ³⁵S cysteine (Amersham) were added. The cells were further incubated for 16 hours before they were harvested by centrifugation and the labelled proteins visualized by SDS-PAGE and autoradiography.

Protein from the medium and cytoplasm of the Sf9 cells was analyzed by SDS-PAGE under non-reducing and reducing conditions. See Figures 8A and 8B, respectively. The medium was dialyzed against 50 mM MES, pH 5.8. Precipitates were obtained after dialysis and resuspended in 100 mM NaCitrate, pH 5.0. The resuspended precipitate was analyzed again by SDS-PAGE and was stained with Coomassie Brilliant Blue. See Figure 9.

The medium supernatant was also diluted 1:10 in 50 mM MES, pH 5.8 and applied to an SP-650M column (1.0 x 6.6 cm, Toyopearl) at a flow rate of 1 ml/min. Protein was eluted with step gradients at 200, 300 and 500 mM NaCl. The VEGF2 was obtained using the elution at 500 mM. The eluate was analyzed by SDS-PAGE in the presence or absence of reducing agent, β-mercaptoethanol and stained by Coomassie Brilliant Blue. See Figure 10.

Example 4

Expression of Recombinant VEGF2 in COS Cells

The expression of plasmid, VEGF2-HA is derived from a vector pcDNA1/Amp (Invitrogen) containing: 1) SV40 origin of replication, 2) ampicillin resistance gene, 3) *E.coli* replication origin, 4) CMV promoter

followed by a polylinker region, an SV40 intron and polyadenylation site. A DNA fragment encoding the entire VEGF2 precursor and a HA tag fused in frame to its 3' end was cloned into the polylinker region of the vector, therefore, the recombinant protein expression is directed under the CMV promoter. The HA tag corresponds to an epitope derived from the influenza hemagglutinin protein as previously described (Wilson *et al.*, *Cell* 37:767 (1984)). The infusion of HA tag to the target protein allows easy detection of the recombinant protein with an antibody that recognizes the HA epitope.

The plasmid construction strategy is described as follows:

The DNA sequence encoding VEGF2, ATCC No. 97149, was constructed by PCR using two primers: the 5' primer (CGC GGA TCC ATG ACT GTA CTC TAC CCA) (SEQ ID NO:14) contains a BamH1 site followed by 18 nucleotides of VEGF2 coding sequence starting from the initiation codon; the 3' sequence (CGC TCT AGA TCA AGC GTA GTC TGG GAC GTC GTA TGG GTA CTC GAG GCT CAT TTG TGG TCT 3') (SEQ ID NO:15) contains complementary sequences to an XbaI site, HA tag, XhoI site, and the last 15 nucleotides of the VEGF2 coding sequence (not including the stop codon). Therefore, the PCR product contains a BamHI site, coding sequence followed by an XhoI restriction endonuclease site and HA tag fused in frame, a translation termination stop codon next to the HA tag, and an XbaI site. The PCR amplified DNA fragment and the vector, pcDNA1/Amp, were digested with BamH1 and XbaI restriction enzyme and ligated. The ligation mixture was transformed into *E. coli* strain SURE (Stratagene Cloning Systems, La Jolla, CA 92037) the transformed culture was plated on ampicillin media plates and resistant colonies were selected. Plasmid DNA was isolated from transformants and examined by restriction analysis for the presence of the correct fragment. For expression of the recombinant VEGF2, COS cells were transfected with the expression vector by DEAE-DEXTRAN method (J. Sambrook, E. Fritsch, T. Maniatis, *Molecular Cloning: A Laboratory Manual*, Cold Spring Laboratory Press, (1989)). The expression of the VEGF2-HA protein was detected by radiolabelling and immunoprecipitation method (E.

Harlow and D. Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, (1988)). Cells were labelled for 8 hours with ^{35}S -cysteine two days post transfection. Culture media was then collected and cells were lysed with detergent (RIPA buffer (150 mM NaCl, 1% NP-40, 0.1% SDS, 1% NP-40, 0.5% DOC, 50mM Tris, pH 7.5) (Wilson *et al.*, *Cell* 37:767 (1984)). Both cell lysate and culture media were precipitated with an HA specific monoclonal antibody. Proteins precipitated were analyzed on 15% SDS-PAGE gels.

Example 5

The Effect of Partially Purified VEGF2 Protein on the Growth of Vascular Endothelial Cells

On day 1, human umbilical vein endothelial cells (HUVEC) were seeded at 2.5×10^4 cells/35 mm dish density in M199 medium containing 4% fetal bovine serum (FBS), 16 units/ml heparin, and 50 units/ml endothelial cell growth supplements (ECGS, Biotechnology, Inc.). On day 2, the medium was replaced with M199 containing 10% FBS, 8 units/ml heparin. VEGF2 protein of SEQ ID NO. 2 minus the initial 45 amino acid residues, (VEGF) and basic FGF (bFGF) were added, at the concentration shown. On days 4 and 6, the medium was replaced. On day 8, cell number was determined with a Coulter Counter (See Figure 12).

Example 6

The Effect of Purified VEGF2 Protein on the Growth of Vascular Endothelial Cells

On day 1, human umbilical vein endothelial cells (HUVEC) were seeded at 2.5×10^4 cells/35 mm dish density in M199 medium containing 4% fetal bovine serum (FBS), 16 units/ml heparin, 50 units/ml endothelial cell growth supplements (ECGS, Biotechnology, Inc.). On day 2, the medium was replaced

with M199 containing 10% FBS, 8 units/ml heparin. Purified VEGF2 protein of SEQ ID NO:2 minus initial 45 amino acid residues was added to the medium at this point. On days 4 and 6, the medium was replaced with fresh medium and supplements. On day 8, cell number was determined with a Coulter Counter (See Figure 13).

Example 7

Expression via Gene Therapy

Fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in tissue-culture medium and separated into small pieces. Small chunks of the tissue are placed on a wet surface of a tissue culture flask, approximately ten pieces are placed in each flask. The flask is turned upside down, closed tight and left at room temperature over night. After 24 hours at room temperature, the flask is inverted and the chunks of tissue remain fixed to the bottom of the flask and fresh media (e.g., Ham's F12 media, with 10% FBS, penicillin and streptomycin, is added. This is then incubated at 37°C for approximately one week. At this time, fresh media is added and subsequently changed every several days. After an additional two weeks in culture, a monolayer of fibroblasts emerge. The monolayer is trypsinized and scaled into larger flasks.

pMV-7 (Kirschmeier, P.T. *et al.*, *DNA* 7:219-225 (1988) flanked by the long terminal repeats of the Moloney murine sarcoma virus, is digested with EcoRI and HindIII and subsequently treated with calf intestinal phosphatase. The linear vector is fractionated on agarose gel and purified, using glass beads.

The cDNA encoding a polypeptide of the present invention is amplified using PCR primers which correspond to the 5' and 3' end sequences respectively. The 5' primer containing an EcoRI site and the 3' primer further includes a HindIII site. Equal quantities of the Moloney murine sarcoma virus linear backbone and

the amplified EcoRI and HindIII fragment are added together, in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The ligation mixture is used to transform bacteria HB101, which are then plated onto agar-containing kanamycin for the purpose of confirming that the vector had the gene of interest properly inserted.

The amphotropic pA317 or GP+am12 packaging cells are grown in tissue culture to confluent density in Dulbecco's Modified Eagles Medium (DMEM) with 10% calf serum (CS), penicillin and streptomycin. The MSV vector containing the gene is then added to the media and the packaging cells are transduced with the vector. The packaging cells now produce infectious viral particles containing the gene (the packaging cells are now referred to as producer cells).

Fresh media is added to the transduced producer cells, and subsequently, the media is harvested from a 10 cm plate of confluent producer cells. The spent media, containing the infectious viral particles, is filtered through a millipore filter to remove detached producer cells and this media is then used to infect fibroblast cells. Media is removed from a sub-confluent plate of fibroblasts and quickly replaced with the media from the producer cells. This media is removed and replaced with fresh media. If the titer of virus is high, then virtually all fibroblasts will be infected and no selection is required. If the titer is very low, then it is necessary to use a retroviral vector that has a selectable marker, such as *neo* or *his*.

The engineered fibroblasts are then injected into the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads. The fibroblasts now produce the protein product.

Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, within the scope of the appended claims, the invention may be practiced otherwise than as particularly described.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: HU, JING-SHAN
ROSEN, CRAIG A.
CAO, LIANG

(ii) TITLE OF INVENTION: VASCULAR ENDOTHELIAL GROWTH FACTOR 2

(iii) NUMBER OF SEQUENCES: 15

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: STERNE, KESSLER, GOLDSTEIN & FOX
(B) STREET: 1100 NEW YORK AVENUE
(C) CITY: WASHINGTON
(D) STATE: DC
(E) COUNTRY: USA
(F) ZIP: 20005

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: TO BE ASSIGNED
(B) FILING DATE: HERewith
(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: US 08/207,550
(B) FILING DATE: 8-MAR-1994
(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: US 08/465,968
(B) FILING DATE: 06-JUN-1995
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: MARKOWICZ, KAREN R.
(B) REGISTRATION NUMBER: 36,351
(C) REFERENCE/DOCKET NUMBER: 1488.1000004

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (202)371-2600
(B) TELEFAX: (202)371-2540

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1674 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: sig_peptide

(B) LOCATION: 12..80

(ix) FEATURE:

(A) NAME/KEY: mat_peptide

(B) LOCATION: 81..1268

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 12..1268

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GTCCTTCAC C	ATG CAC TCG CTG GGC TTC TTC TCT GTG GCG TGT TCT CTG	50
	Met His Ser Leu Gly Phe Phe Ser Val Ala Cys Ser Leu	
	-23 -20 -15	
CTC GCC GCT GCG CTG CTC CCG GGT CCT CGC GAG GCG CCC GCC GCC GCC	98	
Leu Ala Ala Ala Leu Leu Pro Gly Pro Arg Glu Ala Pro Ala Ala Ala		
-10 -5 1 5		
GCC GCC TTC GAG TCC GGA CTC GAC CTC TCG GAC GCG GAG CCC GAC GCG	146	
Ala Ala Phe Glu Ser Gly Leu Asp Leu Ser Asp Ala Glu Pro Asp Ala		
10 15 20		
GGC GAG GCC ACG GCT TAT GCA AGC AAA GAT CTG GAG GAG CAG TTA CGG	194	
Gly Glu Ala Thr Ala Tyr Ala Ser Lys Asp Leu Glu Glu Gln Leu Arg		
25 30 35		
TCT GTG TCC AGT GTA GAT GAA CTC ATG ACT GTA CTC TAC CCA GAA TAT	242	
Ser Val Ser Ser Val Asp Glu Leu Met Thr Val Leu Tyr Pro Glu Tyr		
40 45 50		
TGG AAA ATG TAC AAG TGT CAG CTA AGG AAA GGA GGC TGG CAA CAT AAC	290	
Trp Lys Met Tyr Lys Cys Gln Leu Arg Lys Gly Gly Trp Gln His Asn		
55 60 65 70		
AGA GAA CAG GCC AAC CTC AAC TCA AGG ACA GAA GAG ACT ATA AAA TTT	338	
Arg Glu Gln Ala Asn Leu Asn Ser Arg Thr Glu Glu Thr Ile Lys Phe		
75 80 85		
GCT GCA GCA CAT TAT AAT ACA GAG ATC TTG AAA AGT ATT GAT AAT GAG	386	
Ala Ala Ala His Tyr Asn Thr Glu Ile Leu Lys Ser Ile Asp Asn Glu		
90 95 100		
TGG AGA AAG ACT CAA TGC ATG CCA CGG GAG GTG TGT ATA GAT GTG GGG	434	
Trp Arg Lys Thr Gln Cys Met Pro Arg Glu Val Cys Ile Asp Val Gly		
105 110 115		
AAG GAG TTT GGA GTC GCG ACA AAC ACC TTC TTT AAA CCT CCA TGT GTG	482	
Lys Glu Phe Gly Val Ala Thr Asn Thr Phe Phe Lys Pro Pro Cys Val		
120 125 130		
TCC GTC TAC AGA TGT GGG GGT TGC TGC AAT AGT GAG GGG CTG CAG TGC	530	
Ser Val Tyr Arg Cys Gly Gly Cys Cys Asn Ser Glu Gly Leu Gln Cys		
135 140 145 150		

ATG AAC ACC AGC ACG AGC TAC CTC AGC AAG ACG TTA TTT GAA ATT ACA Met Asn Thr Ser Thr Ser Tyr Leu Ser Lys Thr Leu Phe Glu Ile Thr 155 160 165	578
GTG CCT CTC TCT CAA GGC CCC AAA CCA GTA ACA ATC AGT TTT GCC AAT Val Pro Leu Ser Gln Gly Pro Lys Pro Val Thr Ile Ser Phe Ala Asn 170 175 180	626
CAC ACT TCC TGC CGA TGC ATG TCT AAA CTG GAT GTT TAC AGA CAA GTT His Thr Ser Cys Arg Cys Met Ser Lys Leu Asp Val Tyr Arg Gln Val 185 190 195	674
CAT TCC ATT ATT AGA CGT TCC CTG CCA GCA ACA CTA CCA CAG TGT CAG His Ser Ile Ile Arg Arg Ser Leu Pro Ala Thr Leu Pro Gln Cys Gln 200 205 210	722
GCA GCG AAC AAG ACC TGC CCC ACC AAT TAC ATG TGG AAT AAT CAC ATC Ala Ala Asn Lys Thr Cys Pro Thr Asn Tyr Met Trp Asn Asn His Ile 215 220 225 230	770
TGC AGA TGC CTG GCT CAG GAA GAT TTT ATG TTT TCC TCG GAT GCT GGA Cys Arg Cys Leu Ala Gln Glu Asp Phe Met Phe Ser Ser Asp Ala Gly 235 240 245	818
GAT GAC TCA ACA GAT GGA TTC CAT GAC ATC TGT GGA CCA AAC AAG GAG Asp Asp Ser Thr Asp Gly Phe His Asp Ile Cys Gly Pro Asn Lys Glu 250 255 260	866
CTG GAT GAA GAG ACC TGT CAG TGT GTC TGC AGA GCG GGG CTT CGG CCT Leu Asp Glu Glu Thr Cys Gln Cys Val Cys Arg Ala Gly Leu Arg Pro 265 270 275	914
GCC AGC TGT GGA CCC CAC AAA GAA CTA GAC AGA AAC TCA TGC CAG TGT Ala Ser Cys Gly Pro His Lys Glu Leu Asp Arg Asn Ser Cys Gln Cys 280 285 290	962
GTC TGT AAA AAC AAA CTC TTC CCC AGC CAA TGT GGG GCC AAC CGA GAA Val Cys Lys Asn Lys Leu Phe Pro Ser Gln Cys Gly Ala Asn Arg Glu 295 300 305 310	1010
TTT GAT GAA AAC ACA TGC CAG TGT GTA TGT AAA AGA ACC TGC CCC AGA Phe Asp Glu Asn Thr Cys Gln Cys Val Cys Lys Arg Thr Cys Pro Arg 315 320 325	1058
AAT CAA CCC CTA AAT CCT GGA AAA TGT GCC TGT GAA TGT ACA GAA AGT Asn Gln Pro Leu Asn Pro Gly Lys Cys Ala Cys Glu Cys Thr Glu Ser 330 335 340	1106
CCA CAG AAA TGC TTG TTA AAA GGA AAG AAG TTC CAC CAC CAA ACA TGC Pro Gln Lys Cys Leu Leu Lys Gly Lys Lys Phe His His Gln Thr Cys 345 350 355	1154
AGC TGT TAC AGA CGG CCA TGT ACG AAC CGC CAG AAG GCT TGT GAG CCA Ser Cys Tyr Arg Arg Pro Cys Thr Asn Arg Gln Lys Ala Cys Glu Pro 360 365 370	1202
GGA TTT TCA TAT AGT GAA GAA GTG TGT CGT TGT GTC CCT TCA TAT TGG Gly Phe Ser Tyr Ser Glu Glu Val Cys Arg Cys Val Pro Ser Tyr Trp 375 380 385 390	1250
CAA AGA CCA CAA ATG AGC TAAGATTGTA CTGTTTTCCTA GTTCATCGAT	1298

Gln Arg Pro Gln Met Ser
395

TTTCTATTAT GGAAACTGT GTTGCCACAG TAGAACTGTC TGTGAACAGA GAGACCCTTG 1358
TGGGTCCATG CTAACAAAGA CAAAAGTCTG TCTTTCCTGA ACCATGTGGA TAACTTTACA 1418
GAAATGGACT GGAGCTCATC TGCAAAAGGC CTCTTGTAAG GACTGGTTTT CTGCCAATGA 1478
CCAAACAGCC AAGATTTTCC TCTTGTGATT TCTTTAAAAG AATGACTATA TAATTTATTT 1538
CCACTAAAAA TATTGTTTCT GCATTCATTT TTATAGCAAC AACAAATTGGT AAAACTCACT 1598
GTGATCAATA TTTTATATC ATGCAAAATA TGTTTAAAT AAAATGAAAA TTGTATTTAT 1658
AAAAAAAAAA AAAAAA 1674

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 419 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met	His	Ser	Leu	Gly	Phe	Phe	Ser	Val	Ala	Cys	Ser	Leu	Leu	Ala	Ala	
-23			-20					-15					-10			
Ala	Leu	Leu	Pro	Gly	Pro	Arg	Glu	Ala	Pro	Ala	Ala	Ala	Ala	Ala	Phe	
		-5					1				5					
Glu	Ser	Gly	Leu	Asp	Leu	Ser	Asp	Ala	Glu	Pro	Asp	Ala	Gly	Glu	Ala	
10				15					20					25		
Thr	Ala	Tyr	Ala	Ser	Lys	Asp	Leu	Glu	Glu	Gln	Leu	Arg	Ser	Val	Ser	
			30					35					40			
Ser	Val	Asp	Glu	Leu	Met	Thr	Val	Leu	Tyr	Pro	Glu	Tyr	Trp	Lys	Met	
		45					50						55			
Tyr	Lys	Cys	Gln	Leu	Arg	Lys	Gly	Gly	Trp	Gln	His	Asn	Arg	Glu	Gln	
		60				65						70				
Ala	Asn	Leu	Asn	Ser	Arg	Thr	Glu	Glu	Thr	Ile	Lys	Phe	Ala	Ala	Ala	
	75					80					85					
His	Tyr	Asn	Thr	Glu	Ile	Leu	Lys	Ser	Ile	Asp	Asn	Glu	Trp	Arg	Lys	
90				95					100						105	
Thr	Gln	Cys	Met	Pro	Arg	Glu	Val	Cys	Ile	Asp	Val	Gly	Lys	Glu	Phe	
			110					115					120			
Gly	Val	Ala	Thr	Asn	Thr	Phe	Phe	Lys	Pro	Pro	Cys	Val	Ser	Val	Tyr	
		125					130						135			
Arg	Cys	Gly	Gly	Cys	Cys	Asn	Ser	Glu	Gly	Leu	Gln	Cys	Met	Asn	Thr	
		140				145						150				

Ser Thr Ser Tyr Leu Ser Lys Thr Leu Phe Glu Ile Thr Val Pro Leu
155 160 165

Ser Gln Gly Pro Lys Pro Val Thr Ile Ser Phe Ala Asn His Thr Ser
170 175 180 185

Cys Arg Cys Met Ser Lys Leu Asp Val Tyr Arg Gln Val His Ser Ile
190 195 200

Ile Arg Arg Ser Leu Pro Ala Thr Leu Pro Gln Cys Gln Ala Ala Asn
205 210 215

Lys Thr Cys Pro Thr Asn Tyr Met Trp Asn Asn His Ile Cys Arg Cys
220 225 230

Leu Ala Gln Glu Asp Phe Met Phe Ser Ser Asp Ala Gly Asp Asp Ser
235 240 245

Thr Asp Gly Phe His Asp Ile Cys Gly Pro Asn Lys Glu Leu Asp Glu
250 255 260 265

Glu Thr Cys Gln Cys Val Cys Arg Ala Gly Leu Arg Pro Ala Ser Cys
270 275 280

Gly Pro His Lys Glu Leu Asp Arg Asn Ser Cys Gln Cys Val Cys Lys
285 290 295

Asn Lys Leu Phe Pro Ser Gln Cys Gly Ala Asn Arg Glu Phe Asp Glu
300 305 310

Asn Thr Cys Gln Cys Val Cys Lys Arg Thr Cys Pro Arg Asn Gln Pro
315 320 325

Leu Asn Pro Gly Lys Cys Ala Cys Glu Cys Thr Glu Ser Pro Gln Lys
330 335 340 345

Cys Leu Leu Lys Gly Lys Lys Phe His His Gln Thr Cys Ser Cys Tyr
350 355 360

Arg Arg Pro Cys Thr Asn Arg Gln Lys Ala Cys Glu Pro Gly Phe Ser
365 370 375

Tyr Ser Glu Glu Val Cys Arg Cys Val Pro Ser Tyr Trp Gln Arg Pro
380 385 390

Gln Met Ser
395

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 1526 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (ix) FEATURE:
 - (A) NAME/KEY: sig_peptide

(B) LOCATION: 71..142

(ix) FEATURE:

(A) NAME/KEY: mat_peptide

(B) LOCATION: 143..1120

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 71..1120

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CGAGGCCACG GCTTATGCAA GCAAAGATCT GGAGGAGCAG TTACGGTCTG TGTCCAGTGT	60
AGATGAACTC ATG ACT GTA CTC TAC CCA GAA TAT TGG AAA ATG TAC AAG	109
Met Thr Val Leu Tyr Pro Glu Tyr Trp Lys Met Tyr Lys	
-24 -20 -15	
TGT CAG CTA AGG AAA GGA GGC TGG CAA CAT AAC AGA GAA CAG GCC AAC	157
Cys Gln Leu Arg Lys Gly Gly Trp Gln His Asn Arg Glu Gln Ala Asn	
-10 -5 1 5	
CTC AAC TCA AGG ACA GAA GAG ACT ATA AAA TTT GCT GCA GCA CAT TAT	205
Leu Asn Ser Arg Thr Glu Glu Thr Ile Lys Phe Ala Ala Ala His Tyr	
10 15 20	
AAT ACA GAG ATC TTG AAA AGT ATT GAT AAT GAG TGG AGA AAG ACT CAA	253
Asn Thr Glu Ile Leu Lys Ser Ile Asp Asn Glu Trp Arg Lys Thr Gln	
25 30 35	
TGC ATG CCA CGG GAG GTG TGT ATA GAT GTG GGG AAG GAG TTT GGA GTC	301
Cys Met Pro Arg Glu Val Cys Ile Asp Val Gly Lys Glu Phe Gly Val	
40 45 50	
GCG ACA AAC ACC TTC TTT AAA CCT CCA TGT GTG TCC GTC TAC AGA TGT	349
Ala Thr Asn Thr Phe Phe Lys Pro Pro Cys Val Ser Val Tyr Arg Cys	
55 60 65	
GGG GGT TGC TGC AAT AGT GAG GGG CTG CAG TGC ATG AAC ACC AGC ACG	397
Gly Gly Cys Cys Asn Ser Glu Gly Leu Gln Cys Met Asn Thr Ser Thr	
70 75 80 85	
AGC TAC CTC AGC AAG ACG TTA TTT GAA ATT ACA GTG CCT CTC TCT CAA	445
Ser Tyr Leu Ser Lys Thr Leu Phe Glu Ile Thr Val Pro Leu Ser Gln	
90 95 100	
GGC CCC AAA CCA GTA ACA ATC AGT TTT GCC AAT CAC ACT TCC TGC CGA	493
Gly Pro Lys Pro Val Thr Ile Ser Phe Ala Asn His Thr Ser Cys Arg	
105 110 115	
TGC ATG TCT AAA CTG GAT GTT TAC AGA CAA GTT CAT TCC ATT ATT AGA	541
Cys Met Ser Lys Leu Asp Val Tyr Arg Gln Val His Ser Ile Ile Arg	
120 125 130	
CGT TCC CTG CCA GCA ACA CTA CCA CAG TGT CAG GCA GCG AAC AAG ACC	589
Arg Ser Leu Pro Ala Thr Leu Pro Gln Cys Gln Ala Ala Asn Lys Thr	
135 140 145	
TGC CCC ACC AAT TAC ATG TGG AAT AAT CAC ATC TGC AGA TGC CTG GCT	637
Cys Pro Thr Asn Tyr Met Trp Asn Asn His Ile Cys Arg Cys Leu Ala	

150	155	160	165	
CAG GAA GAT TTT ATG TTT TCC TCG GAT GCT GGA GAT GAC TCA ACA GAT				685
Gln Glu Asp Phe Met Phe Ser Ser Asp Ala Gly Asp Asp Ser Thr Asp	170	175	180	
GGA TTC CAT GAC ATC TGT GGA CCA AAC AAG GAG CTG GAT GAA GAG ACC				733
Gly Phe His Asp Ile Cys Gly Pro Asn Lys Glu Leu Asp Glu Glu Thr	185	190	195	
TGT CAG TGT GTC TGC AGA GCG GGG CTT CGG CCT GCC AGC TGT GGA CCC				781
Cys Gln Cys Val Cys Arg Ala Gly Leu Arg Pro Ala Ser Cys Gly Pro	200	205	210	
CAC AAA GAA CTA GAC AGA AAC TCA TGC CAG TGT GTC TGT AAA AAC AAA				829
His Lys Glu Leu Asp Arg Asn Ser Cys Gln Cys Val Cys Lys Asn Lys	215	220	225	
CTC TTC CCC AGC CAA TGT GGG GCC AAC CGA GAA TTT GAT GAA AAC ACA				877
Leu Phe Pro Ser Gln Cys Gly Ala Asn Arg Glu Phe Asp Glu Asn Thr	230	235	240	245
TGC CAG TGT GTA TGT AAA AGA ACC TGC CCC AGA AAT CAA CCC CTA AAT				925
Cys Gln Cys Val Cys Lys Arg Thr Cys Pro Arg Asn Gln Pro Leu Asn	250	255	260	
CCT GGA AAA TGT GCC TGT GAA TGT ACA GAA AGT CCA CAG AAA TGC TTG				973
Pro Gly Lys Cys Ala Cys Glu Cys Thr Glu Ser Pro Gln Lys Cys Leu	265	270	275	
TTA AAA GGA AAG AAG TTC CAC CAC CAA ACA TGC AGC TGT TAC AGA CGG				1021
Leu Lys Gly Lys Lys Phe His His Gln Thr Cys Ser Cys Tyr Arg Arg	280	285	290	
CCA TGT ACG AAC CGC CAG AAG GCT TGT GAG CCA GGA TTT TCA TAT AGT				1069
Pro Cys Thr Asn Arg Gln Lys Ala Cys Glu Pro Gly Phe Ser Tyr Ser	295	300	305	
GAA GAA GTG TGT CGT TGT GTC CCT TCA TAT TGG CAA AGA CCA CAA ATG				1117
Glu Glu Val Cys Arg Cys Val Pro Ser Tyr Trp Gln Arg Pro Gln Met	310	315	320	325
AGC TAAGATTGTA CTGTTTTCCA GTTCATCGAT TTTCTATTAT GGAAAACTGT				1170
Ser				
GTTGCCACAG TAGAACTGTC TGTGAACAGA GAGACCCTTG TGGGTCCATG CTAACAAAGA				1230
CAAAAGTCTG TCTTTCCTGA ACCATGTGGA TAACTTTACA GAAATGGACT GGAGCTCATC				1290
TGCAAAAGGC CTCTTGTAAG GACTGGTTTT CTGCCAATGA CCAAACAGCC AAGATTTTCC				1350
TCTTGTGATT TCTTTAAAAG AATGACTATA TAATTTATTT CCACTAAAAA TATTGTTTCT				1410
GCATTCATTT TTATAGCAAC AACAATTGGT AAAACTCACT GTGATCAATA TTTTATATC				1470
ATGCAAAATA TGTTTAAAAT AAAATGAAAA TTGTATTTAT AAAAAAAAAA AAAAAA				1526

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 350 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

```

Met Thr Val Leu Tyr Pro Glu Tyr Trp Lys Met Tyr Lys Cys Gln Leu
-24          -20          -15          -10

Arg Lys Gly Gly Trp Gln His Asn Arg Glu Gln Ala Asn Leu Asn Ser
      -5              1              5

Arg Thr Glu Glu Thr Ile Lys Phe Ala Ala Ala His Tyr Asn Thr Glu
      10              15              20

Ile Leu Lys Ser Ile Asp Asn Glu Trp Arg Lys Thr Gln Cys Met Pro
 25          30          35          40

Arg Glu Val Cys Ile Asp Val Gly Lys Glu Phe Gly Val Ala Thr Asn
      45          50          55

Thr Phe Phe Lys Pro Pro Cys Val Ser Val Tyr Arg Cys Gly Gly Cys
      60          65          70

Cys Asn Ser Glu Gly Leu Gln Cys Met Asn Thr Ser Thr Ser Tyr Leu
      75          80          85

Ser Lys Thr Leu Phe Glu Ile Thr Val Pro Leu Ser Gln Gly Pro Lys
  90          95          100

Pro Val Thr Ile Ser Phe Ala Asn His Thr Ser Cys Arg Cys Met Ser
105          110          115          120

Lys Leu Asp Val Tyr Arg Gln Val His Ser Ile Ile Arg Arg Ser Leu
      125          130          135

Pro Ala Thr Leu Pro Gln Cys Gln Ala Ala Asn Lys Thr Cys Pro Thr
      140          145          150

Asn Tyr Met Trp Asn Asn His Ile Cys Arg Cys Leu Ala Gln Glu Asp
      155          160          165

Phe Met Phe Ser Ser Asp Ala Gly Asp Asp Ser Thr Asp Gly Phe His
      170          175          180

Asp Ile Cys Gly Pro Asn Lys Glu Leu Asp Glu Glu Thr Cys Gln Cys
185          190          195          200

Val Cys Arg Ala Gly Leu Arg Pro Ala Ser Cys Gly Pro His Lys Glu
      205          210          215

Leu Asp Arg Asn Ser Cys Gln Cys Val Cys Lys Asn Lys Leu Phe Pro
      220          225          230

Ser Gln Cys Gly Ala Asn Arg Glu Phe Asp Glu Asn Thr Cys Gln Cys
      235          240          245

Val Cys Lys Arg Thr Cys Pro Arg Asn Gln Pro Leu Asn Pro Gly Lys

```

250		255		260											
Cys	Ala	Cys	Glu	Cys	Thr	Glu	Ser	Pro	Gln	Lys	Cys	Leu	Leu	Lys	Gly
265					270					275					280
Lys	Lys	Phe	His	His	Gln	Thr	Cys	Ser	Cys	Tyr	Arg	Arg	Pro	Cys	Thr
			285						290					295	
Asn	Arg	Gln	Lys	Ala	Cys	Glu	Pro	Gly	Phe	Ser	Tyr	Ser	Glu	Glu	Val
		300						305					310		
Cys	Arg	Cys	Val	Pro	Ser	Tyr	Trp	Gln	Arg	Pro	Gln	Met	Ser		
	315						320					325			

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 196 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Met	Arg	Thr	Leu	Ala	Cys	Leu	Leu	Leu	Leu	Gly	Cys	Gly	Tyr	Leu	Ala
1				5					10					15	
His	Val	Leu	Ala	Glu	Glu	Ala	Glu	Ile	Pro	Arg	Glu	Val	Ile	Glu	Arg
		20					25						30		
Leu	Ala	Arg	Ser	Gln	Ile	His	Ser	Ile	Arg	Asp	Leu	Gln	Arg	Leu	Leu
		35					40					45			
Glu	Ile	Asp	Ser	Val	Gly	Ser	Glu	Asp	Ser	Leu	Asp	Thr	Ser	Leu	Arg
	50					55					60				
Ala	His	Gly	Val	His	Ala	Thr	Lys	His	Val	Pro	Glu	Lys	Arg	Pro	Leu
65				70					75						80
Pro	Ile	Arg	Arg	Lys	Arg	Ser	Ile	Glu	Glu	Ala	Val	Pro	Ala	Val	Cys
			85					90						95	
Lys	Thr	Arg	Thr	Val	Ile	Tyr	Glu	Ile	Pro	Arg	Ser	Gln	Val	Asp	Pro
		100					105						110		
Thr	Ser	Ala	Asn	Phe	Leu	Ile	Trp	Pro	Pro	Cys	Val	Glu	Val	Lys	Arg
		115					120					125			
Cys	Thr	Gly	Cys	Cys	Asn	Thr	Ser	Ser	Val	Lys	Cys	Gln	Pro	Ser	Arg
	130					135					140				
Val	His	His	Arg	Ser	Val	Lys	Val	Ala	Lys	Val	Glu	Tyr	Val	Arg	Lys
145					150					155					160
Lys	Pro	Lys	Leu	Lys	Glu	Val	Gln	Val	Arg	Leu	Glu	Glu	His	Leu	Glu
			165					170						175	

Cys Ala Cys Ala Thr Thr Ser Leu Asn Pro Asp Tyr Arg Glu Glu Asp
 180 185 190
 Thr Asp Val Arg
 195

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 241 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met Asn Arg Cys Trp Ala Leu Phe Leu Ser Leu Cys Cys Tyr Leu Arg
 1 5 10 15
 Leu Val Ser Ala Glu Gly Asp Pro Ile Pro Glu Glu Leu Tyr Glu Met
 20 25 30
 Leu Ser Asp His Ser Ile Arg Ser Phe Asp Asp Leu Gln Arg Leu Leu
 35 40 45
 His Gly Asp Pro Gly Glu Glu Asp Gly Ala Glu Leu Asp Leu Asn Met
 50 55 60
 Thr Arg Ser His Ser Gly Gly Glu Leu Glu Ser Leu Ala Arg Gly Arg
 65 70 75 80
 Arg Ser Leu Gly Ser Leu Thr Ile Ala Glu Pro Ala Met Ile Ala Glu
 85 90 95
 Cys Lys Thr Arg Thr Glu Val Phe Glu Ile Ser Arg Arg Leu Ile Asp
 100 105 110
 Arg Thr Asn Ala Asn Phe Leu Val Trp Pro Pro Cys Val Glu Val Gln
 115 120 125
 Arg Cys Ser Gly Cys Cys Asn Asn Arg Asn Val Gln Cys Arg Pro Thr
 130 135 140
 Gln Val Gln Leu Arg Pro Val Gln Val Arg Lys Ile Glu Ile Val Arg
 145 150 155 160
 Lys Lys Pro Ile Phe Lys Lys Ala Thr Val Thr Leu Glu Asp His Leu
 165 170 175
 Ala Cys Lys Cys Glu Thr Val Ala Ala Ala Arg Pro Val Thr Arg Ser
 180 185 190
 Pro Gly Gly Ser Gln Glu Gln Arg Ala Lys Thr Pro Gln Thr Arg Val
 195 200 205
 Thr Ile Arg Thr Val Arg Val Arg Arg Pro Pro Lys Gly Lys His Arg

210		215		220
Lys Phe Lys His Thr His Asp Lys Thr Ala Leu Lys Glu Thr Leu Gly				
225		230		235
				240
Ala				

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 232 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Met	Asn	Phe	Leu	Leu	Ser	Trp	Val	His	Trp	Ser	Leu	Ala	Leu	Leu	Leu
1			5						10				15		
Tyr	Leu	His	His	Ala	Lys	Trp	Ser	Gln	Ala	Ala	Pro	Met	Ala	Glu	Gly
		20						25					30		
Gly	Gly	Gln	Asn	His	His	Glu	Val	Val	Lys	Phe	Met	Asp	Val	Tyr	Gln
		35					40					45			
Arg	Ser	Tyr	Cys	His	Pro	Ile	Glu	Thr	Leu	Val	Asp	Ile	Phe	Gln	Glu
	50					55					60				
Tyr	Pro	Asp	Glu	Ile	Glu	Tyr	Ile	Phe	Lys	Pro	Ser	Cys	Val	Pro	Leu
65				70						75				80	
Met	Arg	Cys	Gly	Gly	Cys	Cys	Asn	Asp	Glu	Gly	Leu	Glu	Cys	Val	Pro
			85						90					95	
Thr	Glu	Glu	Ser	Asn	Ile	Thr	Met	Gln	Ile	Met	Arg	Ile	Lys	Pro	His
			100					105					110		
Gln	Gly	Gln	His	Ile	Gly	Glu	Met	Ser	Phe	Leu	Gln	His	Asn	Lys	Cys
		115					120					125			
Glu	Cys	Arg	Pro	Lys	Lys	Asp	Arg	Ala	Arg	Gln	Glu	Lys	Lys	Ser	Val
	130					135					140				
Arg	Gly	Lys	Gly	Lys	Gly	Gln	Lys	Arg	Lys	Arg	Lys	Lys	Ser	Arg	Tyr
145				150					155						160
Lys	Ser	Trp	Ser	Val	Tyr	Val	Gly	Ala	Arg	Cys	Cys	Leu	Met	Pro	Trp
			165						170					175	
Ser	Leu	Pro	Gly	Pro	His	Pro	Cys	Gly	Pro	Cys	Ser	Glu	Arg	Arg	Lys
			180					185					190		
His	Leu	Phe	Val	Gln	Asp	Pro	Gln	Thr	Cys	Lys	Cys	Ser	Cys	Lys	Asn
		195					200					205			

Thr Asp Ser Arg Cys Lys Ala Arg Gln Leu Glu Leu Asn Glu Arg Thr
210 215 220

Cys Arg Cys Asp Lys Pro Arg Arg
225 230

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 14 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Pro Xaa Cys Val Xaa Xaa Xaa Arg Cys Xaa Gly Cys Cys Asn
1 5 10

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ATGCTTCCGG CTCGTATG

18

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GGGTTTTCCC AGTCACGAC

18

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

CCACATGGTT CAGGAAAGAC A

21

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 50 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

TGTAATACGA CTCACTATAG GGATCCCGCC ATGGAGGCCA CGGCTTATGC

50

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GATCTCTAGA TTAGCTCATT TGTGGTCT

28

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

CGCGGATCCA TGACTGTACT CTACCCA

27

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 60 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

CGCTCTAGAT CAAGCGTAGT CTGGGACGTC GTATGGGTAC TCGAGGCTCA TTTGTGGTCT

60

CGCTCTAGAT CAAGCGTAGT CTGGGACGTC GTATGGGTAC TCGAGGCTCA TTTGTGGTCT

What Is Claimed Is:

1. An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence at least 95% identical to a sequence selected from the group consisting of:

5 (a) a nucleotide sequence encoding a polypeptide comprising amino acids from about -23 to about 396 in SEQ ID NO:2;

(b) a nucleotide sequence encoding a polypeptide comprising amino acids from about - 22 to about 396 in SEQ ID NO:2;

10 (c) a nucleotide sequence encoding a polypeptide comprising amino acids from about 1 to about 396 in SEQ ID NO:2;

(d) a nucleotide sequence encoding a polypeptide comprising amino acids from about -24 to about 326 in SEQ ID NO:4;

(e) a nucleotide sequence encoding a polypeptide comprising amino acids from about - 23 to about 326 in SEQ ID NO:4;

15 (f) a nucleotide sequence encoding a polypeptide comprising amino acids from about 1 to about 326 in SEQ ID NO:4;

(g) a nucleotide sequence encoding a polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit Nos. 97149 or 75698;

20 (h) a nucleotide sequence encoding the mature VEGF2 polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 97149 or 75698; and

(i) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), (c), (d), (e), (f), (g), or (h).

25 2. The nucleic acid molecule of claim 1, wherein said polynucleotide has the nucleotide sequence in SEQ ID NO:1.

3. The nucleic acid molecule of claim 1, wherein said polynucleotide has the nucleotide sequence in SEQ ID NO:3.

4. The nucleic acid molecule of claim 1, wherein said polynucleotide has the nucleotide sequence in SEQ ID NO:1 encoding the VEGF2 polypeptide having the amino acid sequence in SEQ ID NO:2.

5. The nucleic acid molecule of claim 1, wherein said polynucleotide has the nucleotide sequence in SEQ ID NO:3 encoding the VEGF2 polypeptide having the amino acid sequence in SEQ ID NO:4.

6. The nucleic acid molecule of claim 1, wherein said polynucleotide has the nucleotide sequence in SEQ ID NO:1 encoding the mature VEGF2 polypeptide having the amino acid sequence in SEQ ID NO:2.

7. The nucleic acid molecule of claim 1, wherein said polynucleotide has the nucleotide sequence in SEQ ID NO:3 encoding the mature VEGF2 polypeptide having the amino acid sequence in SEQ ID NO:4.

8. The nucleic acid molecule of claim 1, wherein said polynucleotide has the nucleotide sequence of the cDNA clone contained in ATCC Deposit Nos. 97149 or 75698.

9. The nucleic acid molecule of claim 1, wherein said polynucleotide has the nucleotide sequence encoding the VEGF2 polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit Nos. 97149 or 75698.

10. The nucleic acid molecule of claim 1, wherein said polynucleotide has the nucleotide sequence encoding the mature VEGF2 polypeptide having the

amino acid sequence encoded by the cDNA clone contained in ATCC Deposit Nos. 97149 or 75698.

5 11. An isolated nucleic acid molecule comprising a polynucleotide which hybridizes under stringent hybridization conditions to a polynucleotide having a nucleotide sequence identical to a nucleotide sequence in (a), (b), (c), (d), (e), (f), (g), (h), or (i) of claim 1 wherein said polynucleotide which hybridizes does not hybridize under stringent hybridization conditions to a polynucleotide having a nucleotide sequence consisting of only A residues or of only T residues.

10 12. A method for making a recombinant vector comprising inserting an isolated nucleic acid molecule of claim 1 into a vector.

13. A recombinant vector produced by the method of claim 12.

14. A method of making a recombinant host cell comprising introducing the recombinant vector of claim 13 into a host cell.

15 15. A recombinant host cell produced by the method of claim 14.

16. A recombinant method for producing a VEGF2 polypeptide, comprising culturing the recombinant host cell of claim 15 under conditions such that said polypeptide is expressed and recovering said polypeptide.

20 17. An isolated VEGF2 polypeptide having an amino acid sequence at least 95% identical to a sequence selected from the group consisting of:

- (a) amino acids from about -23 to about 396 in SEQ ID NO:2;
- (b) amino acids from about -22 to about 396 in SEQ ID NO:2;
- (c) amino acids from about 1 to about 396 in SEQ ID NO:2;

- (d) amino acids from about -24 to about 326 in SEQ ID NO:4;
- (e) amino acids from about -23 to about 326 in SEQ ID NO:4;
- (f) amino acids from about 1 to about 326 in SEQ ID NO:4;
- (g) the amino acid sequence of the VEGF2 polypeptide having
5 the amino acid sequence encoded by the cDNA clones contained in ATCC
Deposit Nos. 97149 or 75698; and
- (h) the amino acid sequence of the mature VEGF2
polypeptide having the amino acid sequence encoded by the cDNA clone
contained in ATCC Deposit Nos. 97149 or 75698.

10 18. The isolated polypeptide of claim 17, which is produced or
contained in a recombinant host cell.

19. The isolated polypeptide of claim 18, wherein said recombinant
host cell is mammalian.

15 20. An isolated nucleic acid molecule comprising a polynucleotide
encoding a VEGF2 polypeptide wherein, except for at least one conservative
amino acid substitution, said polypeptide has a sequence selected from the group
consisting of:

- (a) a nucleotide sequence encoding a polypeptide comprising
amino acids from about -23 to about 396 in SEQ ID NO:2;
- 20 (b) a nucleotide sequence encoding a polypeptide comprising
amino acids from about - 22 to about 396 in SEQ ID NO:2;
- (c) a nucleotide sequence encoding a polypeptide comprising
amino acids from about 1 to about 396 in SEQ ID NO:2;
- (d) a nucleotide sequence encoding a polypeptide comprising
25 amino acids from about -24 to about 326 in SEQ ID NO:4;
- (e) a nucleotide sequence encoding a polypeptide comprising
amino acids from about - 23 to about 326 in SEQ ID NO:4;

(f) a nucleotide sequence encoding a polypeptide comprising amino acids from about 1 to about 326 in SEQ ID NO:4;

(g) a nucleotide sequence encoding a polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit Nos. 97149 or 75698;

(h) a nucleotide sequence encoding the mature VEGF2 polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 97149 or 75698; and

(i) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), (c), (d), (e), (f), (g), or (h).

21. An isolated VEGF2 polypeptide wherein, except for at least one conservative amino acid substitution, said polypeptide has a sequence selected from the group consisting of:

- (a) amino acids from about -23 to about 396 in SEQ ID NO:2;
- (b) amino acids from about -22 to about 396 in SEQ ID NO:2;
- (c) amino acids from about 1 to about 396 in SEQ ID NO:2;
- (d) amino acids from about -24 to about 326 in SEQ ID NO:4;
- (e) amino acids from about -23 to about 326 in SEQ ID NO:4;
- (f) amino acids from about 1 to about 326 in SEQ ID NO:4;
- (g) the amino acid sequence of the VEGF2 polypeptide having the amino acid sequence encoded by the cDNA clones contained in ATCC Deposit Nos. 97149 or 75698; and

(h) the amino acid sequence of the mature VEGF2 polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit Nos. 97149 or 75698.

22. An antibody against the polypeptide of claim 17.

23. A compound effective as an agonist for the polypeptide of claim 17.

24. A compound effective as an antagonist against the polypeptide of claim 17.

5 25. A pharmaceutical composition comprising the polypeptide of claim 17 and a pharmaceutically acceptable carrier.

26. A method for the treatment of a patient having need of VEGF2 comprising: administering to the patient a therapeutically effective amount of the polypeptide of claim 17.

10 27. The method of claim 26, wherein said therapeutically effective amount of the polypeptide is administered by providing to the patient DNA encoding said polypeptide and expressing said polypeptide *in vivo*.

15 28. A method for the treatment of a patient having need of VEGF2 comprising: administering to the patient a therapeutically effective amount of the compound of claim 23.

29. A method for the treatment of a patient having need to inhibit VEGF2 comprising: administering to the patient a therapeutically effective amount of the antagonist of claim 24.

20 30. A process for diagnosing a disease or a susceptibility to a disease related to expression of the polypeptide of claim 17 comprising:

determining a mutation in the nucleic acid sequence encoding said polypeptide.

31. A diagnostic process comprising:
analyzing for the presence of the polypeptide of claim 17 in a sample derived
from a host.

32. A method for identifying compounds which bind to and activate
or inhibit a receptor for the polypeptide of claim 17 comprising:

contacting a cell expressing on the surface thereof a receptor for the
polypeptide, said receptor being associated with a second component capable of
providing a detectable signal in response to the binding of a compound to said
receptor, with a compound to be screened under conditions to permit binding to
the receptor; and

determining whether the compound binds to and activates or inhibits the
receptor by detecting the presence or absence of a signal generated from the
interaction of the compound with the receptor.

Vascular Endothelial Growth Factor 2

Abstract

Disclosed are human VEGF2 polypeptides, biologically active, diagnostically or therapeutically useful fragments, analogs, or derivatives thereof, and DNA(RNA) encoding such VEGF2 polypeptides. Also provided are procedures for producing such polypeptides by recombinant techniques and antibodies and antagonists against such polypeptides. Such polypeptides may be used therapeutically for stimulating wound healing and for vascular tissue repair. Also provided are methods of using the antibodies and antagonists to inhibit tumor angiogenesis and thus tumor growth, inflammation, diabetic retinopathy, rheumatoid arthritis, and psoriasis.

FIG. 1A

```

1  GTCTTCCACCATGCACCTCGCTGGGCTTCTCTGTGTGGCGTGTTCTCTGCTCCGCTG
  -----+-----+-----+-----+-----+-----+-----+-----+
61  CAGGAAGGTGGTACGTGAGCGACCCGAAGAAGACACCCGACAGAAGACGAGCGCGAC
    M H S L G F F S V A C S L L A A A -
C                                     -

  CGCTGCTCCCGGGTCTCGGAGGCGCCCGCCGCGCGCGCTTCGAGTCCGACTCG
  -----+-----+-----+-----+-----+-----+-----+-----+
61  GCGACGAGGCGCCAGGAGCGCTCCGCGGCGCGCGCGCGGAGAGCTCAGGCGCTGAGC
    L L P G P R E A P A A A A F E S G L D -
C                                     -

  ACCTCTCGGACGCGGAGCCCGGAGCGCGCGGAGGCCACGCTTATGCAAGCAAGATCTGG
  -----+-----+-----+-----+-----+-----+-----+-----+
121  TGGAGAGCTGCGCCTCGGGCTGCGCGCCGCTCCGGTGCCGAATACGTTCTCTAGACC
    L S D A E P D A G E A T A Y A S K D L E -
C                                     -

  AGGAGCAGTTACGGTCTGTGTTCCAGTGTAGATGAATCATGACTGACTCTACCAGAAT
  -----+-----+-----+-----+-----+-----+-----+-----+
181  TCCTCGTCAATGCCAGACACAGGTCAATCTTACTTGAGTACTGACATGAGATGGTCTTA
    E Q L R S V S S V D E L M T V L Y P E Y -
C                                     -

  ATTGAAAATGTACAAGTGTACGCTAAGGAAGAAGGCTGGCAACATAACAGAGAACAGG
  -----+-----+-----+-----+-----+-----+-----+-----+
241  TAACCTTTTACATGTTACACAGTGCATTCCTTTCCTCCGACCGTTGTATTGTCTTGTCC
    W K M Y K C Q L R K G G W Q H N R E Q A -
C                                     -

  CCAACCTCAACTCAAGGACAGAAGAGACTATATAAATTGCTGCAGCACAATTATAATACAG

```

MATCH WITH FIG. 1B

100 100 100 100 100 100 100 100 100 100
 200 200 200 200 200 200 200 200 200 200
 300 300 300 300 300 300 300 300 300 300

FIG. 1

THE AFFAIRS OF THE EAST

NEW YORK TIMES

1 D V G K E F G V A T N T F F K P P C V -

SECRET

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
84

MATCH WITH FIG. 1C

FIG. 1C

GTGCTGTAACAACAACTCTTCCCCAGCCCATGTGGGCCCAACCAGAATTTCATGA
961 -----+-----+-----+-----+-----+-----+-----+
CACAGACATTTTCTTTTGAGAGGGGTCGGTACACCCCGGTTGGCTCTPAACTACTTT
MATCH WITH FIG. 1D 98

FIG. 1D

V C K N K L F P S Ö C G A N R E F D E N -

1021 +-----+ 1080

C T C Q C V C K R T C P R N Q P L N P G K -

-----+-----+-----+-----+-----+
1081

-----+-----+-----+-----+-----+ 1140

C C A C C E C T E S P Q K C L L K G K K F H -

[illegible]


C H Q T C S C Y R R P C T N R Q K A C E P -

[illegible]

C G F S S E E V C R C V P S Y W O R P Q -

AAATGAGCTAAGATTGTA CTGTTTCCAGTTCATCGATTCTATTA TGGAACA CTGCT

MATCH WITH FIG. 1E




```

1  CGAGGCCACGGCTTATGCAAGCAAGATCTGGAGGAGCAGTTACGGTCTGTGTCACACTGT
-----+-----+-----+-----+-----+-----+-----+
71 AGATGAACTCATGACTGTACTCTACCCAGAATATTGAAAATGTACAAGTGTCAAGCTAAG
-----+-----+-----+-----+-----+-----+-----+
      M T V L Y P E Y W K M Y K C Q L R
-----+-----+-----+-----+-----+-----+-----+
121 GAAAGGAGGCTGGCAACATAACAGAGAACAGCCCACTCACTCAAGGACAGAGAGAC
-----+-----+-----+-----+-----+-----+-----+
      K G G W Q H N R E Q A N L N S R T E E T
-----+-----+-----+-----+-----+-----+-----+
181 TATAAAATTGCTGCAGCACATTATAATACAGAGATCTTGAAAAGTATTGATAATGAGTG
-----+-----+-----+-----+-----+-----+-----+
      I K F A A A H Y N T E I L K S I D N E W
-----+-----+-----+-----+-----+-----+-----+
241 GAGAAAGACTCAATGCATGCCACGGAGGTGTGTATAGATGTGGGAAGGAGTTGGAGT
-----+-----+-----+-----+-----+-----+-----+
      R K T Q C M P R E V C I D V G K E F G V
-----+-----+-----+-----+-----+-----+-----+
301 CGCGACAACACCTTCTTTAAACCTCCATGTGTGTCCGTCTACAGATGTGGGGTTCCTG
-----+-----+-----+-----+-----+-----+-----+
      A T N T F F K P P C V S V Y R C G G C C
-----+-----+-----+-----+-----+-----+-----+

```

FIG. 2A

361 CAATAGTGAGGGCTGCAGTGCATGAACACCAGCAGGCTACCTCAGCAAGACGTTAT
 -----+-----+-----+-----+-----+-----+-----+
 N S E G L Q C M N T S T S Y L S K T L F
 421 TGAATTACAGTGCCCTCTCTCTCAAGCCCCCAACCAGTAACAATCAGTTTGGCAATCA
 -----+-----+-----+-----+-----+-----+-----+
 E I T V P L S Q G P K P V T I S F A N H
 481 CACTTCCTGCCGATGCATGTCTAACTGGATGTTTACAGACAAGTTCATTCCATTATTAG
 -----+-----+-----+-----+-----+-----+-----+
 T S C R C M S K L D V Y R Q V H S I I R
 541 ACGTTCCTGCCAGCAACACTACCACAGTGTTCAGCAGCAGCAACAAGACCTGCCCCACCAA
 -----+-----+-----+-----+-----+-----+-----+
 R S L P A T L P Q C Q A A N K T C P T N
 601 TTACATGTGAATAATCACAATCTGCAGATGCCCTGGCTCAGGAAGATTTTATGTTTCCTC
 -----+-----+-----+-----+-----+-----+-----+
 Y M W N N H I C R C L A Q E D F M F S S
 661 GGATGCTGGAGATGACTCAACAGATGGATTCCATGACATCTGTGCAACCAACAAGAGCT
 -----+-----+-----+-----+-----+-----+-----+
 D A G D D S T D G F H D I C G P N K E L

FIG.2B

```

721  GGATGAAGAGACCTGTCAGTGTGCTCGACAGCCGGGCTTCGGCCTGCCAGCTGTGACC
-----+-----+-----+-----+-----+-----+-----+
D E E T C Q C V C R A G L R P A S C G P
-----+-----+-----+-----+-----+-----+-----+
781  CCACAAGAAGCTAGACAGAAACTCATGCCAGTGTGTCTGTA AAAACA AACTCTTCCCCAG
-----+-----+-----+-----+-----+-----+-----+
H K E L D R N S C Q C V C K N K L F P S
-----+-----+-----+-----+-----+-----+-----+
841  CCAATGTGGGCCAACCAGAAATTGATGA AAACACATGCCAGTGTGTATGTAAAGAAC
-----+-----+-----+-----+-----+-----+-----+
Q C G A N R E F D E N T C Q C V C K R T
-----+-----+-----+-----+-----+-----+-----+
901  CTGCCCCAGAAATCAACCCCTAAATCCTGAAAATGTGCCCTGTGAATGTACAGAAAGTCC
-----+-----+-----+-----+-----+-----+-----+
C P R N Q P L N P G K C A C E C T E S P
-----+-----+-----+-----+-----+-----+-----+
961  ACAGAAATGCTTGTAAAGGAAGAAGTTCACACCACCAACAACATGCAGCTGTTACAGACC
-----+-----+-----+-----+-----+-----+-----+
Q K C L L K G K K F H H Q T C S C Y R R
-----+-----+-----+-----+-----+-----+-----+
1021 GCCATGTACGAACCCCGAGAAGGCTTGTGAGCCAGGATTTTCATATAGTGAAGAAGTGTG
-----+-----+-----+-----+-----+-----+-----+
P C T N R Q K A C E P G F S Y S E E V C
-----+-----+-----+-----+-----+-----+-----+

```

FIG. 2C

```

1081 TCCTTGTGTCCTTCATATTGGCAAGACCACAAATGAGCTAAGATTGTA.CTGTTTCCA.
-----+-----+-----+-----+-----+-----+
R C V P S Y W Q R P Q M S
1141 GTTCATCGATTTTCTATTATGGA.AACTGTGTGCCACAGTAGAACTGTCTGTGAACAGA.
-----+-----+-----+-----+-----+-----+
1201 GAGACCCCTGTGGGTCATGCTAACCAAGACAAAGTCTGTCTTCTGAA.CCATGTGA.
-----+-----+-----+-----+-----+-----+
1261 TAACTTTACAGAAATGACTGGAGCTCATCTGCCAAAGCCCTCTGTAAAGACTGGTTT.
-----+-----+-----+-----+-----+-----+
1321 CTGCCAATGACCACAACAGCCAGATTTCCCTCTGTGATTTCTTTAAAGAATGACTATA.
-----+-----+-----+-----+-----+-----+
1381 TAA.TTTATTTCACCTAA.AAATATTGTTCTGCATTCA.TTTTATAGCAACA.CAATTGCT.
-----+-----+-----+-----+-----+-----+
1441 AAAA.CTCACTGTGATCAATATTTTATATCATGCA.AAATATGTTTAA.AAATGA.AAA.
-----+-----+-----+-----+-----+-----+
1501 TTGTATTAT.AAAA.AAAA.AAAA.AAAA.AAAA.AAAA.AAAA.AAAA.AAAA.
-----+-----+-----+-----+-----+-----+

```

FIG.2D

1

50

Pdgfa .MRTIACLLL LGGGYLAHVL AEAEIIPREV IERLARSQIH SIRDLOQLLE
 Pdgtb MNRCWA.LFL SLCCYLRLVS AEGDPIPEEL YEMLSDSIR SFDDLQRLIH
 VegtMNFLL SWHWSLALL LY.....
 Vegf2MTV LYPEYWKMYK CQ..... .LRKGWQH

51

100

Pdgfa IDSVGSEDSL DTSI RAHGVH ATKHVPEKRP LPIRRKRSI.EEAVP
 Pdgtb GDP.GEEDGA ELDLNMTRSH SGELLES... .LARGRSIG SLTIAEPAMI
 Vegt APMAE..... .GGGQ NHHEVVKFMD .VYQR.....
 Vegf2 REQANLNSRT EETIKFAAAH YNTEILKSID NEWRK.....

101

150

Pdgfa AVCKTRTVIX EIPRSQVDP SANFLIMPBC VEVKRCGTGCC NTSSVKCQPS
 Pdgtb AECKTRTEVF EISRRLIDRT NANFLVWPBC VEVQRCSGCC NNRNVOCRPT
 Vegt SYCHPIETLV DIFQEYPDEI .EYIFKPSV PLMRGCGCC NDEGLECVPT
 Vegf2 TQCMPREVCI DVGKEFGVAT ..NTFFKPPC VSVYRCGGCC NSEGLQCMNT

151

200

Pdgfa RVHHRSVKVA KVEYVRKKPK LKEVQVRLEE HLECPAC..... AT.....
 Pdgtb QVQLRPVQVR KIEIVRKPI FKATVTILED HLAACK..... ETVAAARPVT
 Vegt EESNITMQIM RIK.PH..QG QHIGMSFLQ HNKCECRPKK DRARQEKKS
 Vegf2 STSYLSKTLF EIT.VPLSQG PKPVTISFAN HTSCRCMSKL DVYRQVHSII

FIG. 3A

0924 0442 122393

Pd gfa	...	TSLNPD	YREEDTDVR.	250
Pd gfb	RSPGGSQEQR	AKTPQTRVTI	RTVRVRP	PK	GKHKFKHTH	DRTALKETLG
Vegf	RGR.....	..	GKGQKRKRK	KSRYSWSVY	VGARCCIMPW	SLPGPH...
Vegf2	RRSLPATLPQ	CQANKTCPT	NYYMWNHICR	CLAQEDFMFS	SDAGDDSTDG	
	251					
Pd gfa	300
Pd gfb	A.....	
Vegf	...CGP...	
Vegf2	FHDICGNKE	LDEFTCQCVC	RAGLRPASC	G	PHKEL...DR	NSCQCVCXNK
	301					
Pd gfa	350
Pd gfb	
Vegf	..DSRCKARQ	LEINERTCRC	DKPRR.....	
Vegf2	LFPSQCGANR	.EFDENTCQC	VCKRTCPRNQ	PLNPGKACE	CTESPQKLL	
	351					
Pd gfa	398
Pd gfb	
Vegf	
Vegf2	KGKRFHQTC	SCYRRPCTNR	QKACEPGFSY	SEFVCRVPS	YWQRPOMS	


FIG. 3B

PERCENTAGE (%) OF AMINO ACID IDENTITIES BETWEEN
EACH PAIR OF GENES IS SHOWN IN THE
FOLLOWING TABLE

	PDGF α	PDGF β	VEGF	VEGF2
PDGF α				
PDGF β	48.0			
VEGF	20.7	22.7		
VEGF2	23.5	22.4	30.0	


FIG. 4

Expression of VEGF2 mRNA in Human Breast Tumor Cells



- 1. normal breast tissue
- 2. breast tumor tissue
- 3-9. breast tumor cell lines.


FIG. 5



Expression of VEGF2 mRNA in human adult tissues.


FIG. 6

FIG. 7



- Lane 1: 14-C and rainbow M.W. marker
- Lane 2: FGF control
- Lane 3: VEGF2 (M13-reverse & forward primers)
- Lane 4: VEGF2 (M13-reverse & VEGF-F4 primers)
- Lane 5: VEGF2 (M13-reverse & VEGF-F5 primers)


non-reducing gel



Lane M: Marker
Lane 1: vector medium
Lane 2: VEGF2 medium

FIG. 8A

reducing gel



Lane M: Marker
Lane 1: vector cytoplasm
Lane 2: vector medium
Lane 3: VEGF2 cytoplasm
Lane 4: VEGF2 medium

FIG. 8B

FIG. 9




FIG. 10




FIG. 11




FIG. 12





FIG. 13



DECLARATION FOR PATENT APPLICATION

As a below named inventor, I declare that I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

Vascular Endothelial Growth Factor 2

the specification of which was filed on **December 24, 1997** as Application Serial No. **08/999,811**.

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 C.F.R. § 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT international application, which designated at least one country other than the United States listed below, and have also identified below any foreign application for patent or inventor's certificate, or PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s):

Priority Claimed
Yes No

(Number) (Country) (Day/Month/Year Filed)

I hereby claim the benefit under 35 U.S.C. § 119(e) of any United States provisional application(s) listed below.

(Application Serial No.) (Filing Date)


I hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s), or under § 365(b) of any PCT international application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose information that is material to patentability as defined in 37 C.F.R. § 1.56 that became available between the filing date of the prior application and the national or PCT international filing date of this application.

<u>08/207,550</u> (Application Serial No.)	<u>March 8, 1994</u> (Filing Date)	<u>Abandoned</u> (Status: patented, pending, abandoned)
<u>08/465,968</u> (Application Serial No.)	<u>June 6, 1995</u> (Filing Date)	<u>Pending</u> (Status: patented, pending, abandoned)

I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith: A. Anders Brookes (Reg. No. 36,373), James H. Davis (Reg. No. 40,582) and Kenley K. Hoover (Reg. No. 40,302) of Human Genome Sciences, Inc. 9410 Key West Avenue, Rockville, Maryland, 20878.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of first joint inventor: Jing-Shan HU

Inventor's signature:  Date: 7/24/98
Residence: 1247 Lakeside Drive, #3034, Sunnyvale, CA 94086 Citizenship: China
Post Office Address: same as above

Full name of additional joint inventor: Craig A. ROSEN

Inventor's signature: _____ Date: _____
Residence: 22400 Rolling Hill Road, Laytonsville, MD 20882 Citizenship: U.S.
Post Office Address: same as above

Full name of additional joint inventor: Liang CAO

Inventor's signature: _____ Date: _____
Residence: Block 13A, Flat 32D, South Horizons, Hong Kong, China Citizenship: China
Post Office Address: same as above